

# A UNIFIED VIEW OF SOME VERTEX OPERATOR CONSTRUCTIONS

STEPHEN BERMAN<sup>1</sup>, YUN GAO<sup>2</sup>, SHAOBIN TAN<sup>3</sup>

ABSTRACT. We present a general vertex operator construction based on the Fock space for an affine Lie algebras of type  $A$ . This construction allows us to give a unified treatment for both the homogeneous and principle realizations of the affine Lie algebras  $\hat{gl}_N$  as well as for some extended affine Lie algebras coordinatized by certain quantum tori.

## §0. Introduction.

This paper presents a unified view of certain vertex operator constructions for some of the extended affine Lie algebras (EALA's for short) which are coordinatized by certain quantum tori. Recall that for the affine Kac-Moody Lie algebras vertex operator representations were developed in [LW] and [KKLW] for the principal realizations and in [FK],[S] in the homogeneous realization. Our motivation comes from the paper [F1] of I. Frenkel, where he presented a unified construction for both the principal and homogeneous realizations of the affine Lie algebras of type  $A^{(1)}$ . This is accomplished by using the affine algebra  $\hat{gl}_M$  rather than  $\hat{sl}_M$ . Moreover Frenkel used a Clifford algebra structure, which was inherent in his situation, to define a new type of normal ordering which then led to his unified view. The Clifford structure had been studied before in the works [F3,4] and [KP].

The structure theory of EALA's has been developed over the last ten years(see [H-KT], [BGK],[AABGP] and [ABGP]). Roughly speaking these Lie algebras are generalizations of both the affine Kac-Moody Lie algebras and the finite dimensional simple Lie algebras over the complex numbers which admit Laurent like coordinates in a finite number of

---

<sup>1</sup>Department of Mathematics and Statistics, University of Saskatchewan, Saskatoon, Canada S7N 5E6; berman@math.usask.ca

<sup>2</sup>Department of Mathematics and Statistics, York University, Toronto, Canada M3J 1P3; ygao@yorku.ca

<sup>3</sup>Department of Mathematics, Xiamen University, Xiamen, China 361005; tans@jingxian.xmu.edu.cn  
1991 *Mathematics Subject Classification.* 17B10, 17B69, 17B60.

variables. It turns out that algebras of different types admit different types of coordinates. For example, those of type  $A_l$  admit the non-commutative quantum torus as coordinates (see [M],[BGK]). Representations for these Lie algebras over quantum tori have been constructed in [JK], [G-KK], [G1,2,3], [BS], [VV]. When  $l = 1$  there are even Jordan algebras which serve as coordinates of EALAs. The study of representations for this type of Lie algebra has been initiated in [T1]. Perhaps the examples which have attracted the most attention so far are the toroidal algebras which have the commutative associative Laurent polynomials as their coordinates. The toroidal algebras have been studied since the mid 80's both in terms of their structure theory as well as their representation theory (see for example [F2], [MRY], [Y], [W], [BC], [FJW],[T2]). For our purposes we want to mention that vertex operator representations have played a predominant role in much of this work. Indeed, in [G1,2], one finds both homogeneous and principal realizations given for many of the EALA's of type  $A$ . The principal realization for those EALAs was also implicitly given by [G-KK] in studying the so-called  $\Gamma$ -conformal algebras. Our goal in this work is to unify the various approaches and show how they all follow from the same type of approach. Of course, the work in the affine case, namely [F1], shed light on doing this.

Working with a standard type of Fock space we are able to define a general type of vertex operator which depends on a non-zero scalar from  $\mathbb{C}$  and to then compute the commutator of two of these operators. This is presented in the second section of this paper while, in the first section, we give the basics on the Lie algebras, which are all of type  $A$ , which we will later go on to find representations for. Already in Section one it is evident that there is somewhat of a unified picture for these algebras. When we define our vertex operators in Section two the reader will see that we are using a Clifford algebra structure to define the normal ordering we are using, just as was done in [F1]. In the third section we introduce some Lie algebras associated to certain choices of subgroups,  $G$ , of non-zero complex numbers as well as the choice of a positive integer  $M$ . These Lie algebras are spanned by the moments of our vertex operators and hence, by construction, we automatically have a representation for this Lie algebra. We show the representations we have are completely reducible and find the irreducible components. In the fourth and final section we show how certain choices of the group  $G$  and the integer  $M$  lead to representations of the algebras of section one. Thus, we recover both the principal and homogeneous representations for the affine algebras of type  $A$  as well as those for the EALA's studied in [BS], [G1,2], [G-KK]. It is from seeing the various applications in this fourth section that one understands the unification of our treatment. Finally we want to emphasize that this unified treatment would not have been possible without first knowing the particular special cases of this result.

## §1. Preliminaries.

In this section we shall review some of the basics on Lie algebras coordinatized by

quantum tori. We present this from a general point of view which unifies our treatment. For notation we always denote the integers, positive and negative integers respectively by  $\mathbb{Z}$ ,  $\mathbb{Z}_+$ , and  $\mathbb{Z}_-$ .

Let  $\mathfrak{g}$  be any associative  $\mathbb{C}$ -algebra with a symmetric bilinear form  $(\cdot, \cdot): \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{C}$  such that  $(xy, z) = (x, yz)$  for  $x, y, z \in \mathfrak{g}$ . Let  $A = \bigoplus_{\alpha \in \mathbb{Z}^{\nu+1}} A_\alpha$ ,  $\nu \geq 0$ , be any  $\mathbb{Z}^{\nu+1}$ -graded associative algebra such that  $\dim A_\alpha < \infty$  for all  $\alpha \in \mathbb{Z}^{\nu+1}$ . Fix a base  $(x_{i\alpha})_{i \in I_\alpha}$  of  $A_\alpha$ , where  $I_\alpha$  is the index set corresponding to the subspace  $A_\alpha$ . Let  $d_0, d_1, \dots, d_\nu$  be degree derivations of  $A$  such that  $d_i x = \alpha_i x$  for  $x \in A_\alpha$ ,  $i = 0, 1, \dots, \nu$  and  $\alpha = (\alpha_0, \dots, \alpha_\nu) \in \mathbb{Z}^{\nu+1}$ . We define a  $\mathbb{C}$ -linear map  $\phi: A \rightarrow \mathbb{C}$  by linear extension of

$$(1.1) \quad \phi(x_{i\alpha}) = \begin{cases} 1 & \text{if } \alpha = (0, \dots, 0) \\ 0 & \text{otherwise.} \end{cases}$$

for  $i \in I_\alpha$ ,  $\alpha \in \mathbb{Z}^{\nu+1}$ . The tensor product  $\mathfrak{g} \otimes_{\mathbb{C}} A$ , with the canonical product  $(x \otimes a)(y \otimes b) = xy \otimes ab$  for  $x, y \in \mathfrak{g}, a, b \in A$ , is also an associative algebra. Moreover with the commutator product  $[x \otimes a, y \otimes b]_{loop} = (x \otimes a)(y \otimes b) - (y \otimes b)(x \otimes a)$ ,  $\mathfrak{g} \otimes_{\mathbb{C}} A$  forms a  $\mathbb{Z}^{\nu+1}$ -graded Lie algebra. We call this algebra a loop type Lie algebra. Consider the vector space

$$(1.2) \quad \hat{\mathfrak{g}}_A := (\mathfrak{g} \otimes_{\mathbb{C}} A) \oplus \mathcal{C}$$

where  $\mathcal{C} = \bigoplus_{0 \leq i \leq \nu} \mathbb{C} c_i$  is a  $\nu + 1$ -dimensional vector space. There is an alternating bilinear map  $[\cdot, \cdot]: \hat{\mathfrak{g}}_A \times \hat{\mathfrak{g}}_A \rightarrow \hat{\mathfrak{g}}_A$  determined by the conditions:

$$[c_i, \hat{\mathfrak{g}}_A] = 0$$

$$[x \otimes a, y \otimes b] = [x \otimes a, y \otimes b]_{loop} + (x, y) \sum_{0 \leq i \leq \nu} \phi((d_i a)b) c_i$$

for  $x, y \in \mathfrak{g}, a, b \in A$  and  $i = 0, 1, \dots, \nu$ . It is straightforward to check that  $\hat{\mathfrak{g}}_A$  is a Lie algebra. Indeed there is an exact sequence of Lie algebras with canonical maps

$$0 \rightarrow \bigoplus_{0 \leq i \leq \nu} \mathbb{C} c_i \rightarrow \hat{\mathfrak{g}}_A \rightarrow \mathfrak{g} \otimes_{\mathbb{C}} A \rightarrow 0,$$

and so we have that  $\hat{\mathfrak{g}}_A$  is a central extension of the loop type Lie algebra  $\mathfrak{g} \otimes_{\mathbb{C}} A$ .

Let  $M_\infty(\mathbb{C}) = \text{span}_{\mathbb{C}}\{E_{ij} \mid 1 \leq i, j < \infty\}$  be the infinite matrix algebra, where  $E_{ij}$  is the infinite matrix with a 1 in the  $(i, j)$ -entry and zero's elsewhere. We also let  $M_n(\mathbb{C}) = \text{span}_{\mathbb{C}}\{E_{ij} \mid 1 \leq i, j \leq n\}$ . This subspace of  $M_\infty(\mathbb{C})$  for  $n \geq 1$  is isomorphic to the usual matrix algebra of  $n \times n$  matrices with entries in  $\mathbb{C}$ .

Let  $Q = (q_{ij})$  be a  $(\nu + 1) \times (\nu + 1)$  matrix with entries  $q_{ij} \in \mathbb{C}^\times$  satisfying  $q_{ii} = 1$  and  $q_{ij} = q_{ji}^{-1}$  for  $0 \leq i, j \leq \nu$ . The quantum torus associated with the matrix  $Q$  is a unital

associative  $\mathbb{C}$ -algebra  $\mathbb{C}_Q := \mathbb{C}_Q[t_0^{\pm 1}, \dots, t_\nu^{\pm 1}]$  with generators  $t_0^{\pm 1}, \dots, t_\nu^{\pm 1}$  and relations  $t_i t_i^{-1} = t_i^{-1} t_i = 1$ ,  $t_i t_j = q_{ij} t_j t_i$ , for  $0 \leq i, j \leq \nu$ . If  $Q$  is  $2 \times 2$  matrix, so then  $\nu = 1$ , the matrix  $Q = (q_{ij})$  is determined by a single  $q = q_{10}$ . In this case we often simply denote  $\mathbb{C}_Q = \mathbb{C}_Q[t_0^{\pm 1}, t_1^{\pm 1}]$  by  $\mathbb{C}_q$ . Choose the bilinear form on  $M_n(\mathbb{C})$  to be the trace form. Set  $A = \mathbb{C}_Q[t_0^{\pm 1}, \dots, t_\nu^{\pm 1}]$ , with the  $\mathbb{Z}^{\nu+1}$ -gradation  $A = \bigoplus_{\alpha \in \mathbb{Z}^{\nu+1}} A_\alpha$ , where the subspace  $A_\alpha$  is spanned by  $t^\alpha = t_0^{\alpha_0} t_1^{\alpha_1} \dots t_\nu^{\alpha_\nu}$  for  $\alpha = (\alpha_0, \dots, \alpha_\nu) \in \mathbb{Z}^{\nu+1}$ . Define  $\sigma_Q : \mathbb{Z}^{\nu+1} \times \mathbb{Z}^{\nu+1} \rightarrow \mathbb{C}$  by

$$(1.3) \quad \sigma_Q(\alpha, \beta) = \prod_{0 \leq i < j \leq \nu} q_{ji}^{\alpha_j \beta_i}$$

for  $\alpha = (\alpha_0, \dots, \alpha_\nu)$ ,  $\beta = (\beta_0, \dots, \beta_\nu) \in \mathbb{Z}^{\nu+1}$ . Then we have  $t^\alpha t^\beta = \sigma_Q(\alpha, \beta) t^{\alpha+\beta}$ .

The proof of the following Lemma is clear.

**Lemma 1.4.** *Let  $m, n \geq 1$  be integers. Then there is a Lie algebra isomorphism*

$$(M_m(\mathbb{C}) \otimes M_n(\mathbb{C}))_{\mathbb{C}_Q}^\wedge \cong (M_{mn}(\mathbb{C}))_{\mathbb{C}_Q}^\wedge$$

which is given by

$$E_{ij} \otimes E_{kl} \otimes t^\alpha \mapsto E_{(i-1)n+k, (j-1)n+l} \otimes t^\alpha$$

$$c_s \mapsto c_s, \quad s = 0, 1, \dots, \nu$$

for  $\alpha = (\alpha_0, \dots, \alpha_\nu) \in \mathbb{Z}^{\nu+1}$ ,  $1 \leq i, j \leq m$ ,  $1 \leq k, l \leq n$ .

Let  $\hat{\mathcal{L}}_{\mathbb{C}_Q}$  be the Lie subalgebra of  $(M_m(\mathbb{C}) \otimes M_n(\mathbb{C}))_{\mathbb{C}_Q}^\wedge$  generated by elements of the form  $E_{ij} \otimes E_{kl} \otimes t_0^{\alpha_0(n-1)+l-k} t^\alpha$  for  $1 \leq i, j \leq m$ ,  $1 \leq k, l \leq n$  and  $\alpha = (\alpha_0, \dots, \alpha_\nu) \in \mathbb{Z}^{\nu+1}$ . The following result gives the structure of  $\hat{\mathcal{L}}_{\mathbb{C}_Q}$ .

**Proposition 1.5.**

$$\hat{\mathcal{L}}_{\mathbb{C}_Q} \cong (M_m(\mathbb{C}) \otimes M_n(\mathbb{C}))_{\mathbb{C}_{Q^*}}^\wedge$$

where  $\mathbb{C}_Q = \mathbb{C}_Q[t_0^{\pm 1}, \dots, t_\nu^{\pm 1}]$  with  $Q = (q_{ij})$ , and  $\mathbb{C}_{Q^*} = \mathbb{C}_{Q^*}[\tau_0^{\pm 1}, \dots, \tau_\nu^{\pm 1}]$  with  $Q^* = (q_{ij}^*)$  such that  $q_{ij}^* = q_{ij}$  if  $i, j \neq 0$ , and  $q_{ij}^* = q_{ij}^n$  if  $i = 0$  or  $j = 0$ .

Proof. Define a linear map  $f : (M_m(\mathbb{C}) \otimes M_n(\mathbb{C}))_{\mathbb{C}_{Q^*}}^\wedge \rightarrow \hat{\mathcal{L}}_{\mathbb{C}_Q}$  by

$$E_{ij} \otimes E_{kl} \otimes \tau^\alpha \mapsto \left( \prod_{1 \leq s \leq \nu} q_{s0}^{l\alpha_s} \right) E_{ij} \otimes E_{kl} \otimes t_0^{(n-1)\alpha_0+l-k} t^\alpha - k\delta_{ij}\delta_{kl}\delta_{\alpha,0}c_0$$

$$c_0 \mapsto nc_0, \quad c_s \mapsto c_s, \quad s = 1, 2, \dots, \nu$$

for  $\alpha = (\alpha_0, \dots, \alpha_\nu) \in \mathbb{Z}^{\nu+1}$ ,  $1 \leq i, j \leq m$  and  $1 \leq k, l \leq n$ . Let  $\bar{\alpha} = (n\alpha_0 + l - k, \alpha_1, \dots, \alpha_\nu)$ , and  $\bar{\alpha}' = (n\alpha'_0 + l' - k', \alpha'_1, \dots, \alpha'_\nu) \in \mathbb{Z}^{\nu+1}$ . Using the identity

$$(1.6) \quad \sigma_Q(\bar{\alpha}, \bar{\alpha}') = \sigma_{Q^*}(\alpha, \alpha') \prod_{1 \leq j \leq \nu} q_{j0}^{\alpha_j(l' - k')}$$

one can easily check that the map  $f$  is the desired Lie algebra isomorphism.  $\square$

Putting together the two previous results we get the following identification of  $\hat{\mathcal{L}}_{\mathbb{C}_Q}$ .

**Corollary 1.7.**

$$\hat{\mathcal{L}}_{\mathbb{C}Q} \cong (M_{mn}(\mathbb{C}))_{\mathbb{C}Q^*}^\wedge$$

where  $Q$  and  $Q^*$  are given in the previous proposition .  $\square$

Let  $\xi = \xi_n$  be an  $n$ -th primitive root of unity and let  $E, F \in M_n(\mathbb{C})$  be defined by saying

$$(1.8) \quad E = E_{12} + E_{23} + \cdots + E_{n-1,n} + E_{n1}, \quad F = \sum_{i=1}^n E_{ii}(\xi^{i-1}).$$

Then the following fact is well-known.

**Lemma 1.9.** *The set of matrices  $\{F^i E^j\}_{1 \leq i, j \leq n}$ , forms a basis of the matrix algebra  $M_n(\mathbb{C})$  (so a basis of the general linear Lie algebra  $gl_n(\mathbb{C})$ ). Moreover,*

$$(1.10) \quad EF = \xi FE, \quad E^n = F^n = Id_n,$$

and

$$(1.11) \quad E_{ij} = \frac{1}{n} \sum_{k=0}^{n-1} \xi^{k(1-i)} F^k E^{j-i}, \quad F^i E^j = \sum_{l=1}^n \xi^{i(l-1)} E_{l, \overline{l+j}}$$

for  $1 \leq i, j \leq n$ , where, for notation, we are letting  $\bar{l}$  denoted the unique integer,  $l$ , in  $\{1, 2, \dots, n\}$  such that  $\bar{l} = l(\text{mod } n)$ .

Note that

$$\begin{aligned} E_{ij} \otimes E_{kl} \otimes t_0^{\alpha_0(n-1)+l-k} t^\alpha &= \sum_{s=0}^{n-1} \xi^{s(1-k)} E_{ij} \otimes F^s E^{l-k} \otimes t_0^{\alpha_0 n + l - k} t_1^{\alpha_1} \cdots t_\nu^{\alpha_\nu} \\ &= \sum_{s=0}^{n-1} \xi^{s(1-k)} E_{ij} \otimes F^s E^{\alpha'_0} \otimes t_0^{\alpha'_0} t_1^{\alpha_1} \cdots t_\nu^{\alpha_\nu} \end{aligned}$$

where  $\alpha'_0 = \alpha_0 n + l - k$ , for  $1 \leq i, j \leq m$ ,  $1 \leq k, l \leq n$ , and  $\alpha = (\alpha_0, \alpha_1, \dots, \alpha_\nu) \in \mathbb{Z}^{\nu+1}$ . From this one sees that the Lie subalgebra  $\hat{\mathcal{L}}_{\mathbb{C}Q}$  of  $(M_m(\mathbb{C}) \otimes M_n(\mathbb{C}))_{\mathbb{C}Q}^\wedge$  has a basis of the form

$$(1.12) \quad E_{ij} \otimes F^k E^{l_0} \otimes t_0^{l_0} \cdots t_\nu^{l_\nu}, \quad c_0, c_1, \dots, c_\nu$$

where  $1 \leq i, j \leq m$ ,  $0 \leq k \leq n-1$  and  $l_0, l_1, \dots, l_\nu \in \mathbb{Z}$ . Moreover, the commutation relation of  $\hat{\mathcal{L}}_{\mathbb{C}_Q}$  are determined by

$$(1.13) \quad \begin{aligned} & [E_{ij} \otimes F^k E^{\alpha_0} \otimes t^\alpha, E_{i'j'} \otimes F^{k'} E^{\alpha'_0} \otimes t^{\alpha'}] \\ &= \delta_{ji'} \xi^{\alpha_0 k'} \sigma_Q(\alpha, \alpha') E_{ij'} \otimes F^{k+k'} E^{\alpha_0 + \alpha'_0} \otimes t^{\alpha + \alpha'} \\ & - \delta_{j'i} \xi^{\alpha'_0 k} \sigma_Q(\alpha', \alpha) E_{i'j} \otimes F^{k+k'} E^{\alpha_0 + \alpha'_0} \otimes t^{\alpha + \alpha'} + n \delta_{ji'} \delta_{ij'} \delta_{\overline{k+k'}, 0} \delta_{\alpha + \alpha', 0} \xi^{\alpha_0 k'} \sum_{0 \leq s \leq \nu} \alpha_s c_s \end{aligned}$$

for  $1 \leq i, i', j, j' \leq m$ ,  $0 \leq k, k' \leq n-1$ ,  $\alpha = (\alpha_0, \dots, \alpha_\nu)$ ,  $\alpha' = (\alpha'_0, \dots, \alpha'_\nu) \in \mathbb{Z}^{\nu+1}$ , as well as the fact that the elements  $c_0, \dots, c_\nu$  are central in  $\hat{\mathcal{L}}_{\mathbb{C}_Q}$ .

From now on we will identify the Lie algebra  $(M_m(\mathbb{C}) \otimes M_n(\mathbb{C}))_{\mathbb{C}_Q}^\wedge$  with  $(M_{mn}(\mathbb{C}))_{\mathbb{C}_Q}^\wedge$ , and also identify  $\hat{\mathcal{L}}_{\mathbb{C}_Q}$  with  $(M_{mn}(\mathbb{C}))_{\mathbb{C}_{Q^*}}^\wedge$ , where  $Q = (q_{ij})$ ,  $Q^* = (q_{ij}^*)$  and as above  $q_{ij}^* = q_{ij}$  if  $i, j \neq 0$ , and  $q_{ij}^* = q_{ij}^n$  if  $i = 0$  or  $j = 0$ . For simplicity we will write  $a\alpha = (a\alpha_1, \dots, a\alpha_\nu)$  for  $a \in \mathbb{Z}$  and  $\alpha \in \mathbb{Z}^\nu$ , also we will write

$$q_0^\alpha = q_{10}^{\alpha_1} \cdots q_{\nu 0}^{\alpha_\nu}$$

for  $q_0 = (q_{10}, \dots, q_{\nu 0}) \in \mathbb{C}^\nu$ .

The Lie algebra structure (1.13) of  $\hat{\mathcal{L}}_{\mathbb{C}_Q}$  can be described by formal power series identities. For this purpose we let  $z, z_1, z_2$  be formal variables. For  $1 \leq i, j \leq m$ ,  $0 \leq k \leq n-1$  and  $\alpha = (\alpha_1, \dots, \alpha_\nu) \in \mathbb{Z}^\nu$ , we set

$$(1.14) \quad X_{ij}^k(\alpha, z) = \sum_{l \in \mathbb{Z}} (E_{ij} \otimes F^k E^l \otimes t_0^l t_1^{\alpha_1} \cdots t_\nu^{\alpha_\nu}) z^{-l} \in \hat{\mathcal{L}}_{\mathbb{C}_Q}[[z, z^{-1}]],$$

and  $\delta(z) = \sum_{l \in \mathbb{Z}} z^l$ ,  $(D\delta)(z) = \sum_{l \in \mathbb{Z}} l z^l$ . Then the algebra structure of  $\hat{\mathcal{L}}_{\mathbb{C}_Q}$  is described by the following lemma.

**Lemma 1.15.** *Let  $1 \leq i, j, i', j' \leq m$ ,  $0 \leq k, k' \leq n-1$ ,  $\alpha = (\alpha_1, \dots, \alpha_\nu)$ ,  $\alpha' = (\alpha'_1, \dots, \alpha'_\nu) \in \mathbb{Z}^\nu$ . Then the following power series identity is equivalent to (1.13)*

$$(1.16) \quad \begin{aligned} & [X_{ij}^k(\alpha, z_1), X_{i'j'}^{k'}(\alpha', z_2)] = \delta_{ji'} \sigma(\alpha, \alpha') X_{ij'}^{\overline{k+k'}}(\alpha + \alpha', \xi^{-k'} z_1) \delta\left(\frac{\xi^{k'} z_2}{z_1 q_0^\alpha}\right) \\ & - \delta_{j'i} \sigma(\alpha', \alpha) X_{i'j}^{\overline{k+k'}}(\alpha + \alpha', \xi^{-k} z_1) \delta\left(\frac{z_2 q_0^{\alpha'}}{\xi^k z_1}\right) \\ & + n \delta_{ji'} \delta_{ij'} \delta_{\overline{k+k'}, 0} \delta_{\alpha + \alpha', 0} \sigma(\alpha, \alpha') \left\{ (D\delta)\left(\frac{\xi^{k'} z_2}{z_1 q_0^\alpha}\right) c_0 + \delta\left(\frac{\xi^{k'} z_2}{z_1 q_0^\alpha}\right) \sum_{1 \leq s \leq \nu} \alpha_s c_s \right\} \end{aligned}$$

where  $\bar{k} = k(\text{mod } n)$  and  $k \in \{0, 1, \dots, n-1\}$ .

As very special cases, one chooses  $n = 1, \nu = 0$ , then  $\hat{\mathcal{L}}_{\mathbb{C}_Q}$  is just the affine algebra  $\hat{gl}_m(\mathbb{C})$  in the so-called homogeneous picture; while if one chooses  $m = 1, \nu = 0$  then  $\hat{\mathcal{L}}_{\mathbb{C}_Q}$  is the affine algebra  $\hat{gl}_n(\mathbb{C})$  in the so-called principal picture. In these two cases, the identity (1.16) can simply be written as follows

$$(1.17) \quad [X_{ij}^0(z_1), X_{kl}^0(z_2)] = X_{il}^0(z_1)\delta_{jk}\delta\left(\frac{z_2}{z_1}\right) - X_{kj}^0(z_2)\delta_{il}\delta\left(\frac{z_2}{z_1}\right) + \delta_{jk}\delta_{il}(D\delta)\left(\frac{z_2}{z_1}\right)c_0,$$

for  $1 \leq i, j, k, l \leq m$ , for the first of these and

$$(1.18) \quad [X_{11}^i(z_1), X_{11}^j(z_2)] = X_{11}^{\bar{i}+j}(z_2)\delta\left(\frac{\xi^j z_2}{z_1}\right) - X_{11}^{\bar{i}+j}(z_1)\delta\left(\frac{\xi^i z_1}{z_2}\right) + n\delta_{\bar{i}+j,0}(D\delta)\left(\frac{\xi^j z_2}{z_1}\right)c_0.$$

for  $0 \leq i, j \leq n-1, \bar{i} = i(\text{mod } n)$  and  $i \in \{0, 1, \dots, n-1\}$  for the second one.

Moreover if we choose  $n = 1, \nu = 1$ , or  $m = 1, \nu = 1$ , and write  $q_{10} = q$ , then  $\hat{\mathcal{L}}_{\mathbb{C}_Q}$  gives respectively the homogeneous realization of the Lie algebra  $\hat{gl}_m(\mathbb{C}_q)$ , and the principal realization of  $\hat{gl}_n(\mathbb{C}_{q^n})$ . The algebra structure of these two cases can be described as follows

$$(1.19) \quad [X_{ij}^0(r, z_1), X_{kl}^0(s, z_2)] = X_{il}^0(r+s, z_1)\delta_{jk}\delta\left(\frac{z_2}{q^r z_1}\right) - X_{kj}^0(r+s, z_2)\delta_{il}\delta\left(\frac{q^s z_2}{z_1}\right) \\ + \delta_{il}\delta_{jk}\delta_{r+s,0}((D\delta)\left(\frac{z_2}{q^r z_1}\right)c_0 + r\delta\left(\frac{z_2}{q^r z_1}\right)c_1)$$

for  $1 \leq i, j, k, l \leq m$  and  $r, s \in \mathbb{Z}$ , for the first and

$$(1.20) \quad [X_{11}^i(r, z_1), X_{11}^j(s, z_2)] = X_{11}^{\bar{i}+j}(r+s, \xi^{-j} z_1)\delta\left(\frac{\xi^j z_2}{q^r z_1}\right) - X_{11}^{\bar{i}+j}(r+s, \xi^{-i} z_2)\delta\left(\frac{q^s z_2}{\xi^i z_1}\right) \\ + n\delta_{r+s,0}\delta_{\bar{i}+j,0}((D\delta)\left(\frac{\xi^j z_2}{q^r z_1}\right)c_0 + r\delta\left(\frac{\xi^j z_2}{q^r z_1}\right)c_1)$$

for  $1 \leq i, j \leq n, r, s \in \mathbb{Z}$ , and  $\bar{i}+j = i+j(\text{mod } n)$  for the second.

Finally if we choose  $\nu = 1, m, n \geq 1$ , write  $q_{10} = q$ , then  $\hat{\mathcal{L}}_{\mathbb{C}_Q}$  is isomorphic to the affine Lie algebra  $\hat{gl}_{mn}(\mathbb{C}_{q^n})$ , which contains the special cases mentioned above. The algebra structure is as follows.

$$(1.21) \quad [X_{ij}^k(r, z_1), X_{i'j'}^{k'}(r', z_2)] = \delta_{ji'}X_{ij'}^{\bar{k}+k'}(r+r', \xi^{-k'} z_1)\delta\left(\frac{\xi^{k'} z_2}{q^r z_1}\right)$$

$$\begin{aligned}
& -\delta_{j'i} X_{i'j}^{\overline{k+k'}}(r+r', \xi^{-k} z_2) \delta\left(\frac{q^{r'} z_2}{\xi^k z_1}\right) \\
& + n \delta_{ji'} \delta_{ij'} \delta_{\overline{k+k'}, 0} \delta_{r+r', 0} \left\{ (D\delta)\left(\frac{\xi^{k'} z_2}{q^r z_1}\right) c_0 + r \delta\left(\frac{\xi^{k'} z_2}{q^r z_1}\right) c_1 \right\}.
\end{aligned}$$

In subsequent sections we are going to give irreducible representations for a class of Lie algebras which include the Lie algebras mentioned above.

## §2. Fock Space and Vertex Operators.

In this section, we shall define the Fock space we need and construct a family of vertex operators acting on it. Then we go on to derive the commutation relations between these vertex operators in various situations. Some of these commutation relations were implicitly worked out in [G1]. However, we will use the ideas from [F1] to tie a Clifford algebra structure to our vertex operators. This makes our approach very natural and concise.

Let  $\varepsilon_1, \dots, \varepsilon_M$  ( $M \geq 1$ ) be symbols. We form lattices

$$(2.1) \quad \Gamma_M = \oplus_{i=1}^M \mathbb{Z} \varepsilon_i, \quad Q_M = \oplus_{i=1}^{M-1} \mathbb{Z} (\varepsilon_i - \varepsilon_{i+1}),$$

with a symmetric bilinear form  $(\varepsilon_i, \varepsilon_j) = \delta_{ij}$ . We also extend this bilinear form to the  $\mathbb{C}$ -vector space

$$(2.2) \quad H_M := \mathbb{C} \otimes \Gamma_M.$$

For each  $k \in \mathbb{Z}$  we take a copy of  $H_M$  with basis labeled by  $\varepsilon_i(k)$  for  $1 \leq i \leq M, k \in \mathbb{Z}$ . That is,  $\varepsilon_i(k)$ , is to be a copy of  $\varepsilon_i$ . We form a Lie algebra

$$(2.3) \quad \mathcal{H}_M = \text{span}_{\mathbb{C}} \{ \varepsilon_i(k), c | 1 \leq i \leq M, k \in \mathbb{Z} \},$$

with the Lie product

$$(2.4) \quad [\alpha(k), \beta(l)] = k(\alpha, \beta) \delta_{k+l, 0} c,$$

for  $\alpha, \beta \in H_M, k, l \in \mathbb{Z}$ , where  $c$  is a central element. Let

$$(2.5) \quad \mathcal{H}_M^{\pm} = \text{span} \{ \varepsilon_i(k) | k \in \mathbb{Z}_{\pm}, 1 \leq i \leq M \}.$$

Then

$$(2.6) \quad \hat{\mathcal{H}}_M = \mathcal{H}_M^+ + \mathbb{C}c + \mathcal{H}_M^-$$



forms a Heisenberg subalgebra of  $\mathcal{H}_M$ .

Let  $\mathcal{S}(\mathcal{H}_M^-)$  be the symmetric algebra over the abelian algebra  $\mathcal{H}_M^-$  and let

$$(2.7) \quad \mathbb{C}[\Gamma_M] := \bigoplus_{\alpha \in \Gamma_M} \mathbb{C}e^\alpha,$$

be the group algebra over  $\Gamma_M$  twisted by a 2-cocycle so that  $e^\alpha e^\beta = \epsilon(\alpha, \beta)e^{\alpha+\beta}$  for  $\alpha, \beta \in \Gamma_M$ . The cocycle

$$(2.8) \quad \epsilon : \Gamma_M \times \Gamma_M \rightarrow \{\pm 1\},$$

is defined by setting

$$(2.9) \quad \epsilon(\varepsilon_i, \varepsilon_j) = 1 \text{ if } i \leq j, \quad \epsilon(\varepsilon_i, \varepsilon_j) = -1 \text{ if } i > j,$$

and

$$(2.10) \quad \epsilon\left(\sum_i m_i \varepsilon_i, \sum_j n_j \varepsilon_j\right) = \prod_{i,j} (\epsilon(\varepsilon_i, \varepsilon_j))^{m_i n_j},$$

for  $m_i, n_j \in \mathbb{Z}$ . One can easily check the following result.

**Lemma 2.11.**  *$\epsilon$  is bi-multiplicative on  $\Gamma_M$ . Moreover.*

$$\epsilon(\alpha, \alpha) = (-1)^{\frac{(\alpha, \alpha)}{2}}, \quad \epsilon(\alpha, \beta)\epsilon(\beta, \alpha) = (-1)^{(\alpha, \beta)}$$

for  $\alpha, \beta \in Q_M$ .

Now we define the Fock space

$$(2.12) \quad V_M = \mathcal{S}(\mathcal{H}_M^-) \otimes \mathbb{C}[\Gamma_M]$$

which affords representations for both the Lie algebra  $\mathcal{H}_M$  and the group algebra  $\mathbb{C}[\Gamma_M]$  with the following actions:

$$\begin{aligned} \varepsilon_i(k).u \otimes e^\beta &= k\left(\frac{\partial}{\partial \varepsilon_i(-k)}u\right) \otimes e^\beta, \text{ for } k \in \mathbb{Z}_+, \\ \varepsilon_i(k).u \otimes e^\beta &= (\varepsilon_i(k)u) \otimes e^\beta, \text{ for } k \in \mathbb{Z}_-, \\ \varepsilon_i(0).u \otimes e^\beta &= (\varepsilon_i, \beta)u \otimes e^\beta, \\ c.u \otimes e^\beta &= u \otimes e^\beta, \text{ and } e^\alpha.u \otimes e^\beta = \epsilon(\alpha, \beta)u \otimes e^{\alpha+\beta}, \end{aligned}$$

for  $\alpha, \beta \in \Gamma_M$ ,  $1 \leq i \leq M$ , and  $u \in \mathcal{S}(\mathcal{H}_M^-)$ . For  $\alpha \in \Gamma_M$ , we define (we are using the standard notation from [FLM])

$$(2.13) \quad \alpha(z) = \sum_{k \in \mathbb{Z}} \alpha(k)z^{-k} \in (\text{End } V_M)[[z, z^{-1}]]$$

and

$$(2.14) \quad E^\pm(\alpha, z) = \exp\left(\sum_{k \in \mathbb{Z}_\pm} \frac{\alpha(k)}{k} z^{-k}\right) \in (\text{End } V_M)[[z, z^{-1}]].$$

Then the follow lemma is straightforward.

**Lemma 2.15.** For  $\alpha, \beta \in \Gamma_M$ ,  $a, b \in \mathbb{C}^\times := \mathbb{C} \setminus \{0\}$ , we have

$$\begin{aligned} E^\pm(0, az) &= 1, [\alpha(k), E^+(\beta, az)] = 0, \text{ if } k \geq 0, \\ [\alpha(z_1), E^\pm(\beta, az_2)] &= -(\alpha, \beta) E^\pm(\beta, az_2) \sum_{k \in \mathbb{Z}_\mp} \left(\frac{az_2}{z_1}\right)^k, \\ E^\pm(\alpha, az) E^\pm(\beta, bz) &= \exp\left(\sum_{k \in \mathbb{Z}_\pm} \frac{1}{k} (a^{-k} \alpha(k) + b^{-k} \beta(k)) z^{-k}\right), \\ E^+(\alpha, az_1) E^-(\beta, bz_2) &= E^-(\beta, bz_2) E^+(\alpha, az_1) \left(1 - \frac{bz_2}{az_1}\right)^{(\alpha, \beta)}. \end{aligned}$$

Let  $v = \alpha_1(-1)^{k_1} \cdots \alpha_r(-r)^{k_r} \otimes e^\beta \in V_M$ , we define a degree operator  $d_0$  of  $V_M$  by setting

$$(2.16) \quad d_0 v = \left(-\sum_{i=1}^r i k_i - \frac{1}{2}(\beta, \beta)\right) v.$$

If  $a$  is any non-zero complex number we define operators

$$(2.17) \quad z^\alpha .u \otimes e^\beta = z^{(\alpha, \beta)} u \otimes e^\beta, \quad a^\alpha .u \otimes e^\beta = a^{(\alpha, \beta)} u \otimes e^\beta$$

for  $\alpha, \beta \in \Gamma_M$ ,  $u \in \mathcal{S}(\mathcal{H}_M^-)$ . Then  $a^\alpha$  is just the evaluation map, at  $a$ , of the operator  $z^\alpha$ . The following result is well-known.

**Lemma 2.18.**

$$\begin{aligned} [d_0, E^\pm(\alpha, az)] &= -D_z E^\pm(\alpha, az) = \left(\sum_{k \in \mathbb{Z}_\pm} \alpha(k) (az)^{-k}\right) E^\pm(\alpha, az), \\ [\alpha(0), z^\beta] &= 0, \quad z^\alpha e^\beta = z^{(\alpha, \beta)} e^\beta z^\alpha \end{aligned}$$

for  $\alpha, \beta \in \Gamma_M$ ,  $a \in \mathbb{C}^\times$ , where  $D_z = \frac{d}{dz}$ .

We will have need to raise some of the complex numbers which arise in our construction below to various powers and care must be taken with this. We thus set up the notation we use for this now. For any complex number  $a \neq 0$ , there is a unique real number  $\theta \in [0, 2\pi)$  such that  $a = |a|e^{\theta\sqrt{-1}}$ . We define

$$Lna = \theta\sqrt{-1} + \ln |a|$$

Viewing  $\mathbb{C}^\times = \mathbb{C} \setminus \{0\}$  as multiplicative group, we call a subgroup  $G$  of  $\mathbb{C}^\times$  **admissible** if  $G = T \times F$ , where  $T = \langle \xi \rangle$  is a cyclic group of finite order  $|T|$  and  $F = \langle q_j | j \in J \rangle$  is a free abelian group with free generators  $q_j, j \in J$ . For  $a = \xi^{-n_0} q_{i_1}^{n_1} \cdots q_{i_k}^{n_k} \in G$ , where  $n_0, n_1, \dots, n_k \in \mathbb{Z}, i_1, \dots, i_k \in J, 0 \leq n_0 \leq |T| - 1$ . We define

$$(2.19) \quad a^r = e^{r(-n_0 L n \xi + n_1 L n q_{i_1} + \cdots + n_k L n q_{i_k})}$$

for  $r \in \mathbb{C}$ .

Recall the definition of limit of formal power series from [FLM]. Let  $V$  be a vector space over  $\mathbb{C}$ . Let

$$f(z_1, z_2) = \sum_{i,j \in \mathbb{Z}} a_{ij} z_1^i z_2^j \in V[[z_1^{\pm 1}, z_2^{\pm 1}]],$$

we say the limit,  $\lim_{z_2 \rightarrow z_1} f(z_1, z_2)$ , exists if, for any  $l \in \mathbb{Z}$ ,  $a_{i,l-i} = 0$  whenever  $|i| \gg 0$ , and write

$$(2.20) \quad \lim_{z_2 \rightarrow z_1} f(z_1, z_2) = f(z_1, z_1) = \sum_{l \in \mathbb{Z}} \left( \sum_{i \in \mathbb{Z}} a_{i,l-i} \right) z_1^l.$$

For our purposes we need another notion of limit as well. Let  $f(x, z) = \sum_{i \in \mathbb{Z}} C_i(x) z^i$ , where  $C_i(x) = \sum_j c_{ij}(x) v_j \in \mathbb{C}(x) \otimes V$ , and  $j$  runs over a finite set for each fixed  $i \in \mathbb{Z}$ , and  $c_{ij}(x) \in \mathbb{C}(x)$  are complex rational functions. We say the limit,  $\lim_{x \rightarrow a} f(x, z)$ , exists if the function  $c_{ij}(x)$  has a usual limit at the point  $a \in \mathbb{C}$  for all  $i, j \in \mathbb{Z}$ , and write

$$(2.21) \quad \lim_{x \rightarrow a} f(x, z) = f(a, z) = \sum_i \left( \sum_j c_{ij}(a) v_j \right) z^i.$$

**Lemma 2.22.** *For  $1 \leq i \leq M$ , we have*

$$(2.23) \quad \lim_{a \rightarrow 1} \frac{1}{1-a} (a^{-\varepsilon_i} - 1) = \varepsilon_i(0),$$

$$(2.24) \quad \lim_{a \rightarrow 1} \frac{1}{1-a} \left( \sum_{k \in \mathbb{Z}_{\pm}} \frac{\varepsilon_i(k)}{k} (az)^{-k} - \sum_{k \in \mathbb{Z}_{\pm}} \frac{\varepsilon_i(k)}{k} z^{-k} \right) = \sum_{k \in \mathbb{Z}_{\pm}} \varepsilon_i(k) z^{-k}.$$

*Proof.* For any  $v = u \otimes e^{\beta} \in V_M$ , let  $m = -(\varepsilon_i, \beta) \in \mathbb{Z}$ . Then

$$\frac{1}{1-a} (a^{-\varepsilon_i} - 1).v = \frac{a^m - 1}{1-a} v,$$

and

$$\lim_{a \rightarrow 1} \frac{1}{1-a} (a^{-\varepsilon_i} - 1).v = -mv = \varepsilon_i(0).v$$

as required. The second identity is clear.  $\square$

**Corollary 2.25.**

$$\lim_{a \rightarrow 1} \frac{1}{1-a} (E^\pm(\varepsilon_i, az) - E^\pm(\varepsilon_i, z)) = E^\pm(\varepsilon_i, z) \sum_{k \in \mathbb{Z}_\pm} \varepsilon_i(k) z^{-k}.$$

*Proof.* Note that

$$E^\pm(\varepsilon_i, az) - E^\pm(\varepsilon_i, z) = \sum_{l=1}^{\infty} \frac{1}{l!} \left[ \left( \sum_{k \in \mathbb{Z}_\pm} \frac{\varepsilon_i(k)}{k} (az)^{-k} \right)^l - \left( \sum_{k \in \mathbb{Z}_\pm} \frac{\varepsilon_i(k)}{k} z^{-k} \right)^l \right],$$

and  $A^l - B^l = (A - B) \sum_{j=0}^{l-1} A^{l-1-j} B^j$ , we obtain by applying the previous lemma

$$\begin{aligned} & \lim_{a \rightarrow 1} \frac{1}{1-a} (E^\pm(\varepsilon_i, az) - E^\pm(\varepsilon_i, z)) \\ &= \sum_{l=1}^{\infty} \frac{1}{l!} \left[ l \left( \sum_{k \in \mathbb{Z}_\pm} \frac{\varepsilon_i(k)}{k} z^{-k} \right)^{l-1} \right] \sum_{k \in \mathbb{Z}_\pm} \varepsilon_i(k) z^{-k} = E^\pm(\varepsilon_i, z) \sum_{k \in \mathbb{Z}_\pm} \varepsilon_i(k) z^{-k}. \quad \square \end{aligned}$$

**Corollary 2.26.**

$$\lim_{a \rightarrow 1} \frac{1}{1-a} (E^\pm(-\varepsilon_i, z) E^\pm(\varepsilon_i, az) - 1) = \sum_{k \in \mathbb{Z}_\pm} \varepsilon_i(k) z^{-k}.$$

*Proof.* This follows from the fact that

$$E^\pm(-\varepsilon_i, z) E^\pm(\varepsilon_i, az) - 1 = E^\pm(-\varepsilon_i, z) (E^\pm(\varepsilon_i, az) - E^\pm(\varepsilon_i, z)),$$

and the previous corollary.  $\square$

For  $\alpha \in \Gamma_M$ , we define

$$(2.27) \quad X(\alpha, z) = E^-(-\alpha, z) E^+(\alpha, z) e^\alpha z^\alpha z^{(\alpha, \alpha)/2}.$$

We may formally write

$$(2.28) \quad X(\alpha, z) = \sum_{k \in \mathbb{Z} + (\alpha, \alpha)/2} x_k(\alpha) z^{-k},$$

where  $x_k(\alpha) \in \text{End}(V_M)$  for  $k \in \mathbb{Z} + (\alpha, \alpha)/2$ . It is known from [F1] that, if  $(\alpha, \alpha) = 1$ , the operators  $\{x_k(\alpha), x_k(-\alpha) | k \in \mathbb{Z} + \frac{1}{2}\}$  generate a Clifford algebra with the relations

$$(2.29) \quad \{x_k(\alpha), x_{-l}(-\alpha)\} = \delta_{kl}, \quad \{x_k(\alpha), x_l(\alpha)\} = 0, \quad \{x_k(-\alpha), x_l(-\alpha)\} = 0$$

for all  $k, l \in \mathbb{Z} + \frac{1}{2}$ . Related to this Clifford structure, we define the following normal ordering (see [F1] and [G3]):

$$(2.30) \quad : x_k(\varepsilon_i) x_{-l}(-\varepsilon_j) := x_k(\varepsilon_i) x_{-l}(-\varepsilon_j) - \delta_{ij} \delta_{kl} \theta(k)$$

for  $k, l \in \mathbb{Z} + \frac{1}{2}$ ,  $1 \leq i, j \leq M$ , where  $\theta(k) = 0$  if  $k < 0$ ,  $\theta(k) = 1$  if  $k > 0$ . By applying (2.15) and (2.18), one can easily prove the following result

**Lemma 2.31.** For  $1 \leq i, j \leq M$ , and  $a \in \mathbb{C}^\times$ , we have

$$\begin{aligned} & : X(\varepsilon_i, z_1) X(-\varepsilon_j, az_2) : \\ &= \left(1 - \frac{az_2}{z_1}\right)^{-\delta_{ij}} \frac{(az_2)^{\delta_{ij}/2}}{z_1^{\delta_{ij}/2}} \left(\epsilon(\varepsilon_i, \varepsilon_j) z_1^{\frac{(\varepsilon_i, \varepsilon_i - \varepsilon_j)}{2}} \cdot (az_2)^{-\frac{(\varepsilon_j, \varepsilon_i - \varepsilon_j)}{2}}\right) \\ & \cdot e^{\varepsilon_i - \varepsilon_j} z_1^{\varepsilon_i} (az_2)^{-\varepsilon_j} E^-(-\varepsilon_i, z_1) E^-(\varepsilon_j, az_2) E^+(-\varepsilon_i, z_1) E^+(\varepsilon_j, az_2) - \delta_{ij}. \end{aligned}$$

In particular, if  $i \neq j$ , then

$$(2.32) \quad : X(\varepsilon_i, z_1) X(-\varepsilon_j, az_2) := \epsilon(\varepsilon_i, \varepsilon_j) z_1^{\frac{1}{2}} (az_2)^{\frac{1}{2}} e^{\varepsilon_i - \varepsilon_j} z_1^{\varepsilon_i} (az_2)^{-\varepsilon_j} \\ \cdot E^-(-\varepsilon_i, z_1) E^-(\varepsilon_j, az_2) E^+(-\varepsilon_i, z_1) E^+(\varepsilon_j, az_2),$$

and, if  $i = j$ , then

$$(2.33) \quad \left(1 - \frac{az_2}{z_1}\right) : X(\varepsilon_i, z_1) X(-\varepsilon_i, az_2) : \\ = \frac{(az_2)^{\frac{1}{2}}}{z_1^{\frac{1}{2}}} \left(\left(\frac{z_1}{az_2}\right)^{\varepsilon_i} E^-(-\varepsilon_i, z_1) E^-(\varepsilon_i, az_2) E^+(-\varepsilon_i, z_1) E^+(\varepsilon_i, az_2) - 1\right).$$

**Proposition 2.34.** For  $1 \leq i, j \leq M$ , and  $a \in \mathbb{C}^\times$ , we have

$$\begin{aligned} & : X(\varepsilon_i, z) X(-\varepsilon_j, az) : \\ &= \begin{cases} \epsilon(\varepsilon_i, \varepsilon_j) z^{\frac{1}{2}} (az)^{\frac{1}{2}} e^{\varepsilon_i - \varepsilon_j} z^{\varepsilon_i} (az)^{-\varepsilon_j} E^-(-\varepsilon_i, z) E^-(\varepsilon_j, az) E^+(-\varepsilon_i, z) E^+(\varepsilon_j, az) \\ \text{if } i \neq j \\ \varepsilon_i(z) & \text{if } i = j, a = 1 \\ \frac{a^{1/2}}{1-a} (a^{-\varepsilon_i} E^-(-\varepsilon_i, z) E^-(\varepsilon_i, az) E^+(-\varepsilon_i, z) E^+(\varepsilon_i, az) - 1) \\ \text{if } i = j, a \neq 1. \end{cases} \end{aligned}$$

*Proof.* Taking the limit  $z_2 \rightarrow z_1$  in (2.32) and (2.33) gives the first and third identities. The second identity follows from Lemma 2.22, Corollary 2.26 and the third identity by taking the limit  $a \rightarrow 1$ .  $\square$

**Remark 2.35.** Note that the second identity in Proposition 2.34 was given in [F1].

**Definition 2.36.** For  $a \in \mathbb{C}^\times$ ,  $1 \leq i, j \leq M$ , we define  $X_{ij}(a, z) =: X(\varepsilon_i, z) X(-\varepsilon_j, az) :$

Now we can state our main theorem of this section.

**Theorem 2.37.** For  $a, b \in \mathbb{C}^\times$  and  $1 \leq i, j, k, l \leq M$ , we have  
(i) if  $ab \neq 1$ , then

$$(2.38) \quad \begin{aligned} [X_{ij}(a, z_1), X_{kl}(b, z_2)] &= X_{il}(ab, z_1)\delta_{jk}\delta\left(\frac{z_2}{az_1}\right) - X_{kj}(ab, z_2)\delta_{il}\delta\left(\frac{z_1}{bz_2}\right) \\ &\quad + \frac{a^{\frac{1}{2}}b^{\frac{1}{2}}}{1-ab}\delta_{il}\delta_{jk}\left(\delta\left(\frac{z_2}{az_1}\right) - \delta\left(\frac{z_1}{bz_2}\right)\right)c, \end{aligned}$$

(ii) if  $ab = 1$ , then

$$(2.39) \quad [X_{ij}(a, z_1), X_{kl}(b, z_2)] = (X_{il}(1, z_1)\delta_{jk} - X_{kj}(1, z_2)\delta_{il})\delta\left(\frac{z_2}{az_1}\right) + \delta_{il}\delta_{jk}(D\delta)\left(\frac{z_2}{az_1}\right)c.$$

The proof of Theorem 2.37 will be carried out in several steps. In what follows we will freely use the following two lemmas. (2.40) can be found in [FLM] and [K], and (2.43) can be found in [J], [G1, 2] or [BS].

**Lemma 2.40.** Let  $Y(z_1, z_2)$  be a formal power series in  $z_1, z_2$  with coefficients in a vector space, such that  $\lim_{z_2 \rightarrow z_1} f(z_1, z_2)$  exists. Then

$$(2.41) \quad Y(z_1, z_2)\delta\left(\frac{az_1}{z_2}\right) = Y(z_1, az_1)\delta\left(\frac{az_1}{z_2}\right),$$

$$(2.42) \quad Y(z_1, z_2)(D\delta)\left(\frac{z_2}{az_1}\right) = Y(z_1, az_1)(D\delta)\left(\frac{z_2}{az_1}\right) - (D_{z_2}Y(z_1, z_2))\delta\left(\frac{z_2}{az_1}\right),$$

for  $a \in \mathbb{C}^\times$ .

**Lemma 2.43.** Suppose  $a, b \in \mathbb{C}^\times$ . Then

$$\begin{aligned} &\left(1 - \frac{z_2}{az_1}\right)^{-1}\left(1 - \frac{bz_2}{z_1}\right)^{-1} - \frac{az_1}{z_2}\frac{z_1}{bz_2}\left(1 - \frac{z_1}{bz_2}\right)^{-1}\left(1 - \frac{az_1}{z_2}\right)^{-1} \\ &= \begin{cases} (1-ab)^{-1}\frac{az_1}{z_2}\left(\delta\left(\frac{az_1}{z_2}\right) - \delta\left(\frac{z_1}{bz_2}\right)\right) & \text{if } ab \neq 1 \\ \frac{az_1}{z_2}(D\delta)\left(\frac{z_2}{az_1}\right) & \text{if } ab = 1. \end{cases} \end{aligned}$$

Now we divide the proof for Theorem 2.37 into four different cases.

**Case 1.**  $i \neq j, k \neq l$ .

We obtain, by applying (2.15) and (2.18), that

$$\begin{aligned}
(2.44) \quad & [X_{ij}(a, z_1), X_{kl}(b, z_2)] \\
& = \epsilon(\varepsilon_i, \varepsilon_j) \epsilon(\varepsilon_k, \varepsilon_l) a^{\frac{1}{2}} b^{\frac{1}{2}} e^{\varepsilon_i - \varepsilon_j} e^{\varepsilon_k - \varepsilon_l} a^{-\varepsilon_j} b^{-\varepsilon_l} z_1^{\varepsilon_i - \varepsilon_j} z_2^{\varepsilon_k - \varepsilon_l} z_1 z_2 \\
& \quad \cdot E^-( -\varepsilon_i, z_1) E^-( -\varepsilon_k, z_2) E^-(\varepsilon_j, az_1) E^-(\varepsilon_l, bz_2) \\
& \quad \cdot E^+( -\varepsilon_i, z_1) E^+( -\varepsilon_k, z_2) E^+(\varepsilon_j, az_1) E^+(\varepsilon_l, bz_2) P(z_1, z_2)
\end{aligned}$$

where

$$\begin{aligned}
& P(z_1, z_2) \\
& = a^{-(\varepsilon_j, \varepsilon_k - \varepsilon_l)} z_1^{(\varepsilon_i - \varepsilon_j, \varepsilon_k - \varepsilon_l)} \left(1 - \frac{z_2}{z_1}\right)^{\delta_{ik}} \left(1 - \frac{z_2}{az_1}\right)^{-\delta_{jk}} \left(1 - \frac{bz_2}{z_1}\right)^{-\delta_{il}} \left(1 - \frac{bz_2}{az_1}\right)^{\delta_{jl}} \\
& \quad - (-1)^{(\varepsilon_i - \varepsilon_j, \varepsilon_k - \varepsilon_l)} b^{-(\varepsilon_l, \varepsilon_i - \varepsilon_j)} z_2^{(\varepsilon_k - \varepsilon_l, \varepsilon_i - \varepsilon_j)} \left(1 - \frac{z_1}{z_2}\right)^{\delta_{ik}} \left(1 - \frac{z_1}{bz_2}\right)^{-\delta_{il}} \left(1 - \frac{az_1}{z_2}\right)^{-\delta_{jk}} \left(1 - \frac{az_1}{bz_2}\right)^{\delta_{jl}} \\
& = a^{-(\varepsilon_j, \varepsilon_k - \varepsilon_l)} z_1^{(\varepsilon_i - \varepsilon_j, \varepsilon_k - \varepsilon_l)} \left(1 - \frac{z_2}{z_1}\right)^{\delta_{ik}} \left(1 - \frac{bz_2}{az_1}\right)^{\delta_{jl}} \left( \left(1 - \frac{z_2}{az_1}\right)^{-\delta_{jk}} \left(1 - \frac{bz_2}{z_1}\right)^{-\delta_{il}} \right. \\
& \quad \left. - (-1)^{\delta_{il} + \delta_{jk}} \left(\frac{z_2}{az_1}\right)^{-\delta_{jk}} \left(\frac{bz_2}{z_1}\right)^{-\delta_{il}} \left(1 - \frac{az_1}{z_2}\right)^{-\delta_{jk}} \left(1 - \frac{z_1}{bz_2}\right)^{-\delta_{il}} \right).
\end{aligned}$$

Applying Lemma 2.43 we have two subcases.

*Subcase 1.* If  $ab \neq 1$ , then

$$P(z_1, z_2) = \begin{cases} 0 & \text{if } i \neq l, j \neq k \\ \frac{1}{(1-ab)z_1 z_2} (\delta(\frac{z_2}{az_1}) - \delta(\frac{z_1}{bz_2})) & \text{if } i = l, j = k \\ z_1^{-1} \delta(\frac{z_1}{bz_2}) & \text{if } i = l, j \neq k \\ (az_1)^{-1} \delta(\frac{z_2}{az_1}) & \text{if } i \neq l, j = k. \end{cases}$$

Now (2.38) follows from (2.44) and Lemma 2.40.

*Subcase 2.* If  $ab = 1$ , then

$$\begin{aligned}
P(z_1, z_2) & = a^{-(\varepsilon_j, \varepsilon_k - \varepsilon_l)} z_1^{(\varepsilon_i - \varepsilon_j, \varepsilon_k - \varepsilon_l)} \left(1 - \frac{z_2}{z_1}\right)^{\delta_{ik}} \left(1 - \frac{bz_2}{az_1}\right)^{\delta_{jl}} \left( \left(1 - \frac{z_2}{az_1}\right)^{-\delta_{jk} - \delta_{il}} \right. \\
& \quad \left. - (-1)^{\delta_{il} + \delta_{jk}} \left(\frac{z_2}{az_1}\right)^{-\delta_{jk} - \delta_{il}} \left(1 - \frac{az_1}{z_2}\right)^{-\delta_{jk} - \delta_{il}} \right)
\end{aligned}$$

$$= \begin{cases} 0 & \text{if } i \neq l, j \neq k \\ \frac{1}{z_1 z_2} (D\delta)(\frac{z_2}{az_1}) & \text{if } i = l, j = k \\ z_1^{-1} \delta(\frac{z_2}{az_1}) & \text{if } i = l, j \neq k \\ (az_1)^{-1} \delta(\frac{z_2}{az_1}) & \text{if } i \neq l, j = k, \end{cases}$$

which yields (2.39) by applying Lemma 2.40.

**Case 2.**  $i = j$ ,  $k \neq l$ , and  $ab \neq 1$ .

If  $a = 1$ , then

$$X_{ij}(a, z_1) = \varepsilon_i(z_1).$$

Since

$$\begin{aligned} [\varepsilon_i(z_1), e^{\varepsilon_k - \varepsilon_l}] &= (\delta_{ik} - \delta_{il}) e^{\varepsilon_k - \varepsilon_l}, \\ [\varepsilon_i(z_1), E^\pm(-\varepsilon_k, z_2)] &= \sum_{n \in \mathbb{Z}_\pm} \delta_{ik} \left(\frac{z_1}{z_2}\right)^n E^\pm(-\varepsilon_k, z_2), \end{aligned}$$

and

$$[\varepsilon_i(z_1), E^\pm(\varepsilon_l, bz_2)] = - \sum_{n \in \mathbb{Z}_\pm} \delta_{il} \left(\frac{z_1}{bz_2}\right)^n E^\pm(\varepsilon_l, bz_2),$$

we have

$$\begin{aligned} [X_{ij}(a, z_1), X_{kl}(b, z_2)] &= [\varepsilon_i(z_1), X_{kl}(b, z_2)] = X_{kl}(b, z_2) \\ &\cdot \left( \delta_{ik} - \delta_{il} + \delta_{ik} \sum_{n \in \mathbb{Z}_-} \left(\frac{z_1}{z_2}\right)^n + \delta_{ik} \sum_{n \in \mathbb{Z}_+} \left(\frac{z_1}{z_2}\right)^n - \delta_{il} \sum_{n \in \mathbb{Z}_-} \left(\frac{z_1}{bz_2}\right)^n - \delta_{il} \sum_{n \in \mathbb{Z}_+} \left(\frac{z_1}{bz_2}\right)^n \right) \\ &= X_{kl}(b, z_2) \left( \delta_{ik} \delta\left(\frac{z_2}{z_1}\right) - \delta_{il} \delta\left(\frac{z_1}{bz_2}\right) \right) \\ &= X_{kl}(b, z_1) \delta_{jk} \delta\left(\frac{z_2}{z_1}\right) - X_{kj}(b, z_2) \delta_{il} \delta\left(\frac{z_1}{bz_2}\right), \end{aligned}$$

as needed.

If  $a \neq 1$ , then

$$X_{ij}(a, z_1) = \frac{a^{1/2}}{1-a} (a^{-\varepsilon_i} E^-(-\varepsilon_i, z_1) E^-(\varepsilon_i, az_1) E^+(-\varepsilon_i, z_1) E^+(\varepsilon_i, az_1) - 1),$$



Applying (2.15) and (2.18), we get

$$\begin{aligned}
(2.45) \quad & [X_{ij}(a, z_1), X_{kl}(b, z_2)] \\
&= \frac{a^{1/2}}{1-a} b^{\frac{1}{2}} \epsilon(\varepsilon_k, \varepsilon_l) e^{\varepsilon_k - \varepsilon_l} a^{-\varepsilon_i} b^{-\varepsilon_l} z_2^{\varepsilon_k - \varepsilon_l} z_2 \\
&\quad \cdot E^-(\varepsilon_i, z_1) E^-(\varepsilon_i, az_1) E^-(\varepsilon_k, z_2) E^-(\varepsilon_l, bz_2) \\
&\quad \cdot E^+(\varepsilon_i, z_1) E^+(\varepsilon_i, az_1) E^+(\varepsilon_k, z_2) E^+(\varepsilon_l, bz_2) Q(z_1, z_2),
\end{aligned}$$

where

$$\begin{aligned}
Q(z_1, z_2) &= a^{-(\varepsilon_j, \varepsilon_k - \varepsilon_l)} \left(1 - \frac{z_2}{z_1}\right)^{\delta_{ik}} \left(1 - \frac{z_2}{az_1}\right)^{-\delta_{ik}} \left(1 - \frac{bz_2}{z_1}\right)^{-\delta_{il}} \left(1 - \frac{bz_2}{az_1}\right)^{\delta_{il}} \\
&\quad - \left(1 - \frac{z_1}{z_2}\right)^{\delta_{ik}} \left(1 - \frac{z_1}{bz_2}\right)^{-\delta_{il}} \left(1 - \frac{az_1}{z_2}\right)^{-\delta_{ik}} \left(1 - \frac{az_1}{bz_2}\right)^{\delta_{il}} \\
&= a^{\delta_{il} - \delta_{ik}} \left(1 - \frac{z_2}{z_1}\right)^{\delta_{ik}} \left(1 - \frac{bz_2}{az_1}\right)^{\delta_{il}} \left( \left(1 - \frac{z_2}{az_1}\right)^{-\delta_{ik}} \left(1 - \frac{bz_2}{z_1}\right)^{-\delta_{il}} \right. \\
&\quad \left. - (-1)^{\delta_{ik} + \delta_{il}} \left(\frac{az_1}{z_2}\right)^{\delta_{ik}} \left(\frac{z_1}{bz_2}\right)^{\delta_{il}} \left(1 - \frac{z_1}{bz_2}\right)^{-\delta_{il}} \left(1 - \frac{az_1}{z_2}\right)^{-\delta_{ik}} \right) \\
&= \begin{cases} 0 & \text{if } i \neq k, i \neq l \\ \frac{1-a}{a} \delta\left(\frac{z_2}{az_1}\right) & \text{if } i = k, i \neq l \\ (a-1) \delta\left(\frac{z_1}{bz_2}\right) & \text{if } i \neq k, i = l \end{cases}
\end{aligned}$$

We thus have (2.38) by applying Lemma 2.40.

**Case 3.**  $i = j$ ,  $k \neq l$ , and  $ab = 1$ .

As we did in Case 2, we treat the case  $a = 1$  and  $a \neq 1$  separately. First for  $a = 1$  (so  $b = 1$ ), then

$$X_{ij}(a, z_1) = \varepsilon_i(z_1),$$

$$X_{kl}(b, z_2) = \epsilon(\varepsilon_k, \varepsilon_l) e^{\varepsilon_k - \varepsilon_l} z_2^{\varepsilon_k - \varepsilon_l} E^-(\varepsilon_k, z_2) E^-(\varepsilon_l, z_2) E^+(\varepsilon_k, z_2) E^+(\varepsilon_l, z_2),$$

and we have

$$\begin{aligned}
& [X_{ij}(a, z_1), X_{kl}(b, z_2)] = [\varepsilon_i(z_1), X_{kl}(1, z_2)] \\
&= X_{kl}(1, z_2) \left( \delta_{ik} \delta\left(\frac{z_2}{z_1}\right) - \delta_{il} \delta\left(\frac{z_2}{z_1}\right) \right) \\
&= (X_{il}(1, z_1) \delta_{jk} - X_{kj}(1, z_2) \delta_{il}) \delta\left(\frac{z_2}{z_1}\right),
\end{aligned}$$

as expected.

Next assume that  $a \neq 1$ . By applying similar arguments as in Case 2, we have

$$\begin{aligned}
(2.46) \quad & [X_{ij}(a, z_1), X_{kl}(b, z_2)] \\
&= \frac{1}{1-a} \epsilon(\varepsilon_k, \varepsilon_l) e^{\varepsilon_k - \varepsilon_l} a^{-\varepsilon_i} a^{\varepsilon_l} z_2^{\varepsilon_k - \varepsilon_l} z_2 \\
&\quad \cdot E^-(-\varepsilon_i, z_1) E^-(\varepsilon_i, az_1) E^-(-\varepsilon_k, z_2) E^-(\varepsilon_l, bz_2) \\
&\quad \cdot E^+(-\varepsilon_i, z_1) E^+(\varepsilon_i, az_1) E^+(-\varepsilon_k, z_2) E^+(\varepsilon_l, bz_2) R(z_1, z_2)
\end{aligned}$$

where

$$\begin{aligned}
R(z_1, z_2) &= a^{\delta_{il} - \delta_{ik}} \left(1 - \frac{z_2}{z_1}\right)^{\delta_{ik}} \left(1 - \frac{bz_2}{az_1}\right)^{\delta_{il}} \left(1 - \frac{z_2}{az_1}\right)^{-\delta_{il} - \delta_{ik}} \\
&\quad - (-1)^{\delta_{il} + \delta_{ik}} \left(\frac{az_1}{z_2}\right)^{\delta_{il} + \delta_{ik}} \left(1 - \frac{az_1}{z_2}\right)^{-\delta_{il} - \delta_{ik}} \\
&= \begin{cases} 0 & \text{if } i \neq k, i \neq l \\ (b-1)\delta\left(\frac{z_2}{az_1}\right) & \text{if } i = k, i \neq l \\ (a-1)\delta\left(\frac{z_2}{az_1}\right) & \text{if } i \neq k, i = l. \end{cases}
\end{aligned}$$

Substituting this into (2.46) and applying Lemma 2.40, we get (2.39).

**Case 4.**  $i = j, k = l$ .

Note that

$$\begin{aligned}
& X_{ij}(a, z_1) \\
&= \begin{cases} \frac{a^{1/2}}{1-a} (a^{-\varepsilon_i} E^-(-\varepsilon_i, z_1) E^-(\varepsilon_i, az_1) E^+(-\varepsilon_i, z_1) E^+(\varepsilon_i, az_1) - 1) & \text{if } i = j, a \neq 1 \\ \varepsilon_i(z_1) & \text{if } i = j, a = 1 \end{cases} \\
& X_{kl}(b, z_2) \\
&= \begin{cases} \frac{b^{1/2}}{1-b} (b^{-\varepsilon_k} E^-(-\varepsilon_k, z_2) E^-(\varepsilon_k, bz_2) E^+(-\varepsilon_k, z_2) E^+(\varepsilon_k, bz_2) - 1) & \text{if } k = l, b \neq 1 \\ \varepsilon_k(z_2) & \text{if } k = l, b = 1. \end{cases}
\end{aligned}$$

We consider three subcases. First we assume that  $a = b = 1$ , then

$$[X_{ij}(a, z_1), X_{kl}(b, z_2)] = [\varepsilon_i(z_1), \varepsilon_k(z_2)] = \delta_{ik}(D\delta)\left(\frac{z_2}{z_1}\right),$$

as desired.

Next we assume  $a = 1$ ,  $b \neq 1$ , then

$$\begin{aligned}
& [X_{ij}(a, z_1), X_{kl}(b, z_2)] = [\varepsilon_i(z_1), X_{kl}(b, z_2)] \\
&= \frac{b^{1/2}}{1-b} b^{-\varepsilon_k} E^-(-\varepsilon_k, z_2) E^-(\varepsilon_k, bz_2) E^+(-\varepsilon_k, z_2) E^+(\varepsilon_k, bz_2) \\
&\quad \cdot \left( \delta_{ik} \sum_{n \in \mathbb{Z}_-} \left(\frac{z_2}{z_1}\right)^n + \delta_{ik} \sum_{n \in \mathbb{Z}_+} \left(\frac{z_2}{z_1}\right)^n - \delta_{ik} \sum_{n \in \mathbb{Z}_-} \left(\frac{z_1}{bz_2}\right)^n - \delta_{ik} \sum_{n \in \mathbb{Z}_+} \left(\frac{z_1}{bz_2}\right)^n \right) \\
&= \frac{b^{1/2}}{1-b} b^{-\varepsilon_k} E^-(-\varepsilon_k, z_2) E^-(\varepsilon_k, bz_2) E^+(-\varepsilon_k, z_2) E^+(\varepsilon_k, bz_2) \delta_{ik} \left( \delta\left(\frac{z_2}{z_1}\right) - \delta\left(\frac{z_1}{bz_2}\right) \right) \\
&= X_{il}(ab, z_1) \delta_{jk} \delta\left(\frac{z_2}{az_1}\right) - X_{kj}(ab, z_2) \delta_{il} \delta\left(\frac{z_1}{bz_2}\right) + \frac{a^{\frac{1}{2}} b^{\frac{1}{2}}}{1-ab} \delta_{il} \delta_{jk} \left( \delta\left(\frac{z_2}{az_1}\right) - \delta\left(\frac{z_1}{bz_2}\right) \right),
\end{aligned}$$

as required.

Finally we assume  $a \neq 1$ ,  $b \neq 1$ , then

$$\begin{aligned}
(2.47) \quad & [X_{ij}(a, z_1), X_{kl}(b, z_2)] \\
&= \frac{a^{1/2}}{1-a} \frac{b^{1/2}}{1-b} a^{-\varepsilon_i} b^{-\varepsilon_k} E^-(-\varepsilon_i, z_1) E^-(-\varepsilon_k, z_2) E^-(\varepsilon_i, az_1) \\
&\quad \cdot E^-(\varepsilon_k, bz_2) E^+(-\varepsilon_i, z_1) E^+(-\varepsilon_k, z_2) E^+(\varepsilon_i, az_1) E^+(\varepsilon_k, bz_2) S(z_1, z_2),
\end{aligned}$$

where

$$\begin{aligned}
& S(z_1, z_2) = \left(1 - \frac{z_2}{z_1}\right)^{\delta_{ik}} \left(1 - \frac{bz_2}{z_1}\right)^{-\delta_{ik}} \left(1 - \frac{z_2}{az_1}\right)^{-\delta_{ik}} \left(1 - \frac{bz_2}{az_1}\right)^{\delta_{ik}} \\
&\quad - \left(1 - \frac{z_1}{z_2}\right)^{\delta_{ik}} \left(1 - \frac{z_1}{bz_2}\right)^{-\delta_{ik}} \left(1 - \frac{az_1}{z_2}\right)^{-\delta_{ik}} \left(1 - \frac{az_1}{bz_2}\right)^{\delta_{ik}} \\
&= \left(1 - \frac{z_2}{z_1}\right)^{\delta_{ik}} \left(1 - \frac{bz_2}{az_1}\right)^{\delta_{ik}} \left( \left(1 - \frac{bz_2}{z_1}\right)^{-\delta_{ik}} \left(1 - \frac{z_2}{az_1}\right)^{-\delta_{ik}} \right. \\
&\quad \left. - \left(\frac{az_1}{z_2}\right)^{\delta_{ik}} \left(\frac{z_1}{bz_2}\right)^{\delta_{ik}} \left(1 - \frac{z_1}{bz_2}\right)^{-\delta_{ik}} \left(1 - \frac{az_1}{z_2}\right)^{-\delta_{ik}} \right) \\
&= \begin{cases} \left(1 - \frac{z_2}{z_1}\right) \left(1 - \frac{bz_2}{az_1}\right) \frac{az_1}{z_2} (D\delta)\left(\frac{z_2}{az_1}\right) & \text{if } i = k, ab = 1 \\ 0 & \text{if } i \neq k \\ \frac{1}{1-ab} \left(1 - \frac{z_2}{z_1}\right) \left(1 - \frac{bz_2}{az_1}\right) \frac{az_1}{z_2} \left(\delta\left(\frac{z_2}{az_1}\right) - \delta\left(\frac{z_1}{bz_2}\right)\right) & \text{if } i = k, ab \neq 1. \end{cases}
\end{aligned}$$

Substitute  $S(z_1, z_2)$  back into (2.47) and apply Lemma 2.40 to get (2.38) and (2.39). This now completes the proof of Theorem 2.37.

**Remark 2.48** If  $ab \neq 1$ , we have

$$\delta\left(\frac{z_2}{az_1}\right) - \delta\left(\frac{z_1}{bz_2}\right) = \sum_{m \in \mathbb{Z}} (1 - (ab)^m) \left(\frac{z_2}{az_1}\right)^m,$$

thus

$$\begin{aligned} \frac{a^{\frac{1}{2}} b^{\frac{1}{2}}}{1 - ab} \delta_{il} \delta_{jk} \left( \delta\left(\frac{z_2}{az_1}\right) - \delta\left(\frac{z_1}{bz_2}\right) \right) &= a^{\frac{1}{2}} b^{\frac{1}{2}} \delta_{il} \delta_{jk} \sum_{m \in \mathbb{Z}} \frac{1 - (ab)^m}{1 - ab} \left(\frac{z_2}{az_1}\right)^m \\ &= a^{\frac{1}{2}} b^{\frac{1}{2}} \delta_{il} \delta_{jk} \left( \sum_{m \in \mathbb{Z}_+} \left( \sum_{s=0}^{m-1} (ab)^s \right) \left(\frac{z_2}{az_1}\right)^m + \sum_{m \in \mathbb{Z}_-} \left( - \sum_{s=1}^{-m} (ab)^{-s} \right) \left(\frac{z_2}{az_1}\right)^m \right), \end{aligned}$$

which gives

$$\begin{aligned} \lim_{b \rightarrow a^{-1}} \frac{a^{\frac{1}{2}} b^{\frac{1}{2}}}{1 - ab} \delta_{il} \delta_{jk} \left( \delta\left(\frac{z_2}{az_1}\right) - \delta\left(\frac{z_1}{bz_2}\right) \right) \\ = \delta_{il} \delta_{jk} \left( \sum_{m \in \mathbb{Z}_+} m \left(\frac{z_2}{az_1}\right)^m + \sum_{m \in \mathbb{Z}_-} m \left(\frac{z_2}{az_1}\right)^m \right) = \delta_{il} \delta_{jk} (D\delta) \left(\frac{z_2}{az_1}\right). \end{aligned}$$

This indicates that the second identity of Theorem 2.37 can be obtained from the first one by taking the limit as  $b \rightarrow a^{-1}$ .

### §3. Lie Algebras and Representations.

In this section we are going to define a class of Lie algebras from our vertex operators which will correspond to admissible subgroups of  $\mathbb{C}^\times$ . Indeed for some choice of the admissible group  $G$  and positive integer  $M$ , the Lie algebra  $\mathcal{G}(G, M)$  (defined below) of operators, which act on the Fock space  $V_M$ , will give realizations of some infinite dimensional Lie algebras studied in Section 1. This will include the affine algebra  $\widehat{gl}_M(\mathbb{C})$  in both the principal and homogeneous pictures as well as some Lie algebras with quantum torus coordinates. Towards this end, we first introduce some new notation for the vertex operators constructed in the proceeding section.

**Definition 3.1.** For  $a, b \in \mathbb{C}^\times$ ,  $1 \leq i, j \leq M$ , we set  $X_{ij}(a, b, z) := X_{ij}(a^{-1}b, az)$ , and write

$$(3.2) \quad X_{ij}(a, b, z) = \sum_{k \in \mathbb{Z}} x_{ij}(k, a, b) z^{-k}$$

where  $x_{ij}(k, a, b) \in \text{End} V_M$ .

With this notation Theorem 2.37 can be re-written as follows.

**Theorem 3.3.** Let  $a_1, a_2, b_1, b_2 \in \mathbb{C}^\times$ , and  $1 \leq i, j, k, l \leq M$ . We have

(i) if  $a_1 a_2 \neq b_1 b_2$ , then

$$\begin{aligned} & [X_{ij}(a_1, b_1, z_1), X_{kl}(a_2, b_2, z_2)] \\ &= X_{il}(a_1, \frac{b_1 b_2}{a_2}, z_1) \delta_{jk} \delta(\frac{a_2 z_2}{b_1 z_1}) - X_{kj}(a_2, \frac{b_1 b_2}{a_1}, z_2) \delta_{il} \delta(\frac{a_1 z_1}{b_2 z_2}) \\ &+ \frac{(a_1^{-1} b_1)^{\frac{1}{2}} (a_2^{-1} b_2)^{frac{1}{2}}}{1 - a_1^{-1} b_1 a_2^{-1} b_2} \delta_{il} \delta_{jk} (\delta(\frac{a_2 z_2}{b_1 z_1}) - \delta(\frac{a_1 z_1}{b_2 z_2})) c, \end{aligned}$$

(ii) if  $a_1 a_2 = b_1 b_2$ , then

$$\begin{aligned} & [X_{ij}(a_1, b_1, z_1), X_{kl}(a_2, b_2, z_2)] \\ &= X_{il}(a_1, \frac{b_1 b_2}{a_2}, z_1) \delta_{jk} \delta(\frac{a_2 z_2}{b_1 z_1}) - X_{kj}(a_2, \frac{b_1 b_2}{a_1}, z_2) \delta_{il} \delta(\frac{a_1 z_1}{b_2 z_2}) + \delta_{il} \delta_{jk} (D\delta)(\frac{a_2 z_2}{b_1 z_1}) c. \end{aligned}$$

Fix an integer  $M \geq 1$  and an admissible subgroup  $G$  of  $\mathbb{C}^\times$ . Let  $\mathcal{G}(G, M)$  be the vector space spanned by  $c$  and all of the coefficients of the vertex operators  $X_{ij}(a, b, z)$  for all  $1 \leq i, j \leq M$ , and  $a, b \in G$ . Then we have the following result.

**Theorem 3.4.**  $\mathcal{G}(G, M)$  forms a Lie algebra of operators acting on the Fock space  $V_M$ . Moreover

$$V_M = \oplus_{k \in \mathbb{Z}} V_M^{(k)}$$

where  $V_M^{(k)} = e^{k\varepsilon_M + Q_M} \otimes \mathcal{S}(\mathcal{H}_M^-)$ , and  $V_M^{(k)}$  is an irreducible  $\mathcal{G}(G, M)$ -module.

*Proof.* It is obvious from Theorem 3.3 that  $\mathcal{G}(G, M)$  is a Lie algebra and that  $V_M^{(k)}$  is a  $\mathcal{G}(G, M)$ -module. To see it is irreducible, we note that the Heisenberg algebra  $\widehat{\mathcal{H}}_M \subset \mathcal{H}_M$ , and  $\mathcal{H}_M$  is spanned by the coefficient operators of the vertex operators  $X_{ii}(1, 1, z)$  for  $1 \leq i \leq M$ . This then implies that, if  $W$  is a non-zero submodule of  $V_M^{(k)}$ , we can choose a non-zero element of the form  $v = e^{k\varepsilon_M + \alpha} \otimes 1 \in W$  for some  $\alpha \in Q_M$ . Moreover, it is easy to check that

$$x_{ij}(n_{ij} - 1, 1, 1).v = \epsilon(\varepsilon_i, \varepsilon_j) \epsilon(\varepsilon_i - \varepsilon_j, k\varepsilon_M + \alpha) e^{k\varepsilon_M + \alpha + \varepsilon_i - \varepsilon_j}$$

for all  $1 \leq i \neq j \leq M$ , where  $n_{ij} = (\varepsilon_j - \varepsilon_i, k\varepsilon_M + \alpha) \in \mathbb{Z}$ . Therefore  $e^{k\varepsilon_M + \beta} \otimes 1 \in W$  for any  $\beta \in Q_M$ . This thus gives  $W = V_M^{(k)}$  as needed.  $\square$

**Remark 3.5** Note that the coefficients of the vertex operators  $X_{ij}(a, z)$  and  $X_{ij}(a, bz)$  for any given  $a, b \in G$  span the same space. Thus  $\mathcal{G}(G, M)$  is spanned by  $c$  and the

coefficients of the operators  $X_{ij}(a, z)$  for  $1 \leq i, j \leq M, a \in G$ . Therefore, it follows from Theorem 4.25 in [G1] that  $\mathcal{G}(G, M)$  is an affinization of the Lie algebra  $gl_M(\mathcal{R}[t, t^{-1}; \tau])$ , where  $\mathcal{R} = \mathbb{C}[G]$  is the group algebra and  $\mathcal{R}[t, t^{-1}; \tau]$  is the skew Laurent polynomial ring.

Recall definition (2.19). We extend the cocycle map  $\epsilon : H_M \times H_M \rightarrow \{\mathbb{C}^\times\}$  by defining

$$(3.6) \quad \epsilon\left(\sum r_i \varepsilon_i, \sum s_i \varepsilon_i\right) = \prod_{i,j} (\epsilon(\varepsilon_i, \varepsilon_j))^{r_i s_j}$$

for  $r_i, s_i \in \mathbb{C}$ . It is obvious that

$$\epsilon(\alpha + \beta, \gamma) = \epsilon(\alpha, \gamma)\epsilon(\beta, \gamma), \quad \epsilon(\alpha, \beta + \gamma) = \epsilon(\alpha, \beta)\epsilon(\alpha, \gamma)$$

for  $\alpha, \beta, \gamma \in H_M$ . Moreover, if we restrict  $\epsilon$  to  $\Gamma_M \times \Gamma_M$ , then  $\epsilon$  gives us the 2-cocycle defined in previous section.

Let  $\alpha_i = \varepsilon_i - \varepsilon_{i+1}$ ,  $i = 1, \dots, M-1$ , and  $\alpha_M = \varepsilon_1 + \dots + \varepsilon_M$ . Then  $Q_M = \bigoplus_{i=1}^{M-1} \mathbb{Z}\alpha_i$ , and  $\Gamma_M = \bigoplus_{i=1}^M \mathbb{Z}\alpha_i$ . Let  $Q_M^0 = \{\alpha \in \mathbb{C} \otimes_{\mathbb{Z}} Q_M \mid (\alpha, Q_M) \in \mathbb{Z}\}$  be the dual of the lattice  $Q_M$ , and set

$$(3.7) \quad L_M^0 = \{\alpha \in H_M = \mathbb{C} \otimes_{\mathbb{Z}} \Gamma_M \mid (\alpha, Q_M) \in \mathbb{Z}\}.$$

Then  $L_M^0 = Q_M^0 \oplus \mathbb{C}\alpha_M$ . Let  $I = L_M^0/Q_M$ , then we have a  $Q_M$ -coset decomposition of  $L_M^0$

$$(3.8) \quad L_M^0 = \bigoplus_{i \in I} (\lambda_i + Q_M)$$

for some  $\lambda_i \in L_M^0$ .

**Proposition 3.9.** *The Lie algebra of operators,  $\mathcal{G}(G, M)$ , acts on the space  $V_M^{(\lambda_i)} = \mathcal{S}(\mathcal{H}_M^-) \otimes \mathbb{C}[Q_M + \lambda_i]$ , and  $V_M^{(\lambda_i)}$  affords an irreducible representation of  $\mathcal{G}(G, M)$  for  $i \in I = L_M^0/Q_M$ . Moreover  $V_M^{(\lambda_i)} \cong V_M^{(\lambda_j)}$  if and only if  $i = j$ .*

*Proof.* For  $u \otimes e^{\alpha + \lambda_i} \in V_M^{(\lambda_i)}$ ,  $u \in \mathcal{S}(\mathcal{H}_M^-)$ ,  $\alpha \in Q_M$ , we note that

$$z^\beta \cdot (u \otimes e^{\alpha + \lambda_i}) = z^{(\beta, \alpha + \lambda_i)} u \otimes e^{\alpha + \lambda_i}, \quad a^\gamma \cdot (u \otimes e^{\alpha + \lambda_i}) = a^{(\gamma, \alpha + \lambda_i)} u \otimes e^{\alpha + \lambda_i}$$

$$e^\beta \cdot (u \otimes e^{\alpha + \lambda_i}) = \epsilon(\beta, \alpha + \lambda_i) u \otimes e^{\alpha + \beta + \lambda_i}$$

where  $(\beta, \alpha + \lambda_i) \in \mathbb{Z}$ ,  $(\gamma, \alpha + \lambda_i) \in \mathbb{C}$ ,  $\epsilon(\beta, \alpha + \lambda_i) \in \mathbb{C}^\times$  for  $\beta \in Q_M$ ,  $\gamma \in \Gamma_M$  and  $a \in G$ . This implies that  $X_{ij}(a, b, z) \cdot (u \otimes e^{\alpha + \lambda_i}) \in V_M^{(\lambda_i)}[[z, z^{-1}]]$ , and so the Lie algebra  $\mathcal{G}(G, M)$  acts on the space  $V_M^{(\lambda_i)}$ . The irreducibility of  $V_M^{(\lambda_i)}$ , for  $i \in I$ , follows from a similar argument as in Theorem 3.4. The last part of this proposition is clear.  $\square$

#### §4. Applications.

In this section we assume the admissible subgroup  $G$  has the form  $G = T \times F \subset \mathbb{C}^\times$ , where  $T = \langle \xi \rangle$  is generated by a root of unity  $\xi$ , and  $F$  is a free group with a finite number generators. First, let  $G = \{1\}$ , and  $M \geq 2$  be any integer. Then the Lie algebra  $\mathcal{G}(G, M)$  is generated by the coefficients of the vertex operators  $X_{ij}(1, 1, z)$  for all  $1 \leq i, j \leq M$ . Moreover from Theorem 3.3, we see that

$$(4.1) \quad [X_{ij}(1, 1, z_1), X_{kl}(1, 1, z_2)] \\ = X_{il}(1, 1, z_1) \delta_{jk} \delta\left(\frac{z_2}{z_1}\right) - X_{kj}(1, 1, z_2) \delta_{il} \delta\left(\frac{z_1}{z_2}\right) + \delta_{il} \delta_{jk} (D\delta)\left(\frac{z_2}{z_1}\right)$$

for  $1 \leq i, j \leq M$ . Comparing (4.1) with (1.17), we obtain the following result which was originally due to [F1], see also [FK] and [S].

**Corollary 4.2.** *Let  $G = \{1\}$  and  $M \geq 2$ . Then  $\mathcal{G}(G, M)$  gives a representation of the affine algebra  $\widehat{gl}_M(\mathbb{C})$  in the homogeneous picture on the Fock space  $V_M$ , and the representation is given by the mapping:*

$$E_{ij} \otimes t_0^k \mapsto x_{ij}(k, 1, 1), \\ c_0 \mapsto c$$

for  $1 \leq i, j \leq M$  and  $k \in \mathbb{Z}$ .

Next we choose  $G$  to be a cyclic group of order  $N \geq 2$  with generator  $\xi = \xi_N$ , and take  $M = 1$ . Note that

$$X_{11}(\xi^i, \xi^j, z) = X_{11}(\xi^{i-j-1}, \xi^{-1}, \xi^{j+1} z)$$

for  $0 \leq i, j \leq N-1$ . This implies that the Lie algebra  $\mathcal{G}(G, M)$  is generated by the coefficients of the vertex operators  $X_{11}(\xi^{i-1}, \xi^{-1}, z)$  for  $0 \leq i \leq N-1$ . From Theorem 3.3 we have

$$[X_{11}(\xi^{i-1}, \xi^{-1}, z_1), X_{11}(\xi^{j-1}, \xi^{-1}, z_2)] \\ = \begin{cases} X_{11}(\xi^{i-1}, \xi^{-j-1}, z_1) \delta\left(\frac{\xi^j z_2}{z_1}\right) - X_{11}(\xi^{j-1}, \xi^{-i-1}, z_2) \delta\left(\frac{\xi^i z_1}{z_2}\right) + (D\delta)\left(\frac{\xi^j z_2}{z_1}\right) c \\ \quad \text{if } i+j = 0(\text{mod } N) \\ X_{11}(\xi^{i-1}, \xi^{-j-1}, z_1) \delta\left(\frac{\xi^j z_2}{z_1}\right) - X_{11}(\xi^{j-1}, \xi^{-i-1}, z_2) \delta\left(\frac{\xi^i z_1}{z_2}\right) + \\ \quad \frac{e^{-\frac{i+j}{2} L n \xi}}{\xi^{i+j-1}} \left( \delta\left(\frac{\xi^j z_2}{z_1}\right) - \delta\left(\frac{\xi^i z_1}{z_2}\right) \right) \quad \text{if } i+j \neq 0(\text{mod } N). \end{cases}$$

Recalling the definition of  $X_{ij}(a, b, z)$ , we have

$$X_{11}(\xi^{i-1}, \xi^{-j-1}, z_1) \delta\left(\frac{\xi^j z_2}{z_1}\right) = X_{11}(\xi^{i-1}, \xi^{-j-1}, \xi^j z_2) \delta\left(\frac{\xi^j z_2}{z_1}\right)$$

$$= X_{11}(\xi^{i+j-1}, \xi^{-1}, z_2) \delta\left(\frac{\xi^j z_2}{z_1}\right),$$

while

$$\begin{aligned} X_{11}(\xi^{j-1}, \xi^{-i-1}, z_2) \delta\left(\frac{\xi^i z_1}{z_2}\right) &= X_{11}(\xi^{i-1}, \xi^{-j-1}, \xi^j z_1) \delta\left(\frac{\xi^i z_1}{z_2}\right) \\ &= X_{11}(\xi^{i+j-1}, \xi^{-1}, z_1) \delta\left(\frac{\xi^i z_1}{z_2}\right) \end{aligned}$$

for  $0 \leq i, j \leq N-1$ . Therefore we get

$$(4.3) \quad [X_{11}(\xi^{i-1}, \xi^{-1}, z_1), X_{11}(\xi^{j-1}, \xi^{-1}, z_2)] = \begin{cases} X_{11}(\xi^{i+j-1}, \xi^{-1}, z_2) \delta\left(\frac{\xi^j z_2}{z_1}\right) - X_{11}(\xi^{i+j-1}, \xi^{-1}, z_1) \delta\left(\frac{\xi^i z_1}{z_2}\right) + (D\delta)\left(\frac{\xi^j z_2}{z_1}\right)c & \text{if } i+j = 0(\text{mod } N) \\ X_{11}(\xi^{i+j-1}, \xi^{-1}, z_2) \delta\left(\frac{\xi^j z_2}{z_1}\right) - X_{11}(\xi^{i+j-1}, \xi^{-1}, z_1) \delta\left(\frac{\xi^i z_1}{z_2}\right) + \frac{e^{-\frac{i+j}{2}Ln\xi}}{\xi^{i+j-1}} (\delta\left(\frac{\xi^j z_2}{z_1}\right) - \delta\left(\frac{\xi^i z_1}{z_2}\right)) & \text{if } i+j \neq 0(\text{mod } N)c. \end{cases}$$

Comparing this with the identity (1.18), we obtain the following result which was originally due to [F1] and [KKLW].

**Corollary 4.4.** *Let  $M = 1$ , and let  $G$  be the group generated by  $\xi$ , where  $\xi$  is a  $N$ -th primitive root of unity for  $N \geq 2$ . Then the Lie algebra  $\mathcal{G}(G, M)$  gives a representation of the affine algebra  $\widehat{gl}_N(\mathbb{C})$  on the Fock space  $V_1$  in the principal picture. The representation is given by the mapping*

$$\begin{aligned} F^i E^k \otimes t_0^k &\mapsto \begin{cases} x_{11}(k, \xi^{i-1}, \xi^{-1}) + \frac{e^{\frac{i}{2}Ln\xi}}{\xi^{i-1}} \delta_{k0}c & \text{if } 1 \leq i \leq N-1, \\ x_{11}(k, \xi^{-1}, \xi^{-1}) & \text{if } i = 0 \end{cases}, \\ c_0 &\mapsto \frac{c}{N} \end{aligned}$$

for  $k \in \mathbb{Z}$ .

Next we choose  $M \geq 2$  and  $G = \langle q \rangle$  where  $q \neq 0$  is not a root of unity. Note that  $X_{ij}(a, b, z) = X_{ij}(1, a^{-1}b, az)$  for  $a, b \in G$ . We see that the Lie algebra  $\mathcal{G}(G, M)$  is generated by the coefficients of the vertex operators of the form  $X_{ij}(1, q^r, z)$  for all  $r \in \mathbb{Z}$  and  $1 \leq i, j \leq M$ . We apply Theorem 3.3 to obtain

(i) if  $r + s = 0$ , then

$$(4.5) \quad [X_{ij}(1, q^r, z_1), X_{kl}(1, q^s, z_2)]$$



$$= X_{il}(1, q^{r+s}, z_1) \delta_{jk} \delta\left(\frac{z_2}{q^r z_1}\right) - X_{kj}(1, q^{r+s}, z_2) \delta_{il} \delta\left(\frac{z_1}{q^s z_2}\right) + \delta_{il} \delta_{jk} (D\delta)\left(\frac{z_2}{q^r z_1}\right) c,$$

(ii) if  $r + s \neq 0$ , then

$$(4.6) \quad \begin{aligned} & [X_{ij}(1, q^r, z_1), X_{kl}(1, q^s, z_2)] \\ &= X_{il}(1, q^{r+s}, z_1) \delta_{jk} \delta\left(\frac{z_2}{q^r z_1}\right) - X_{kj}(1, q^{r+s}, z_2) \delta_{il} \delta\left(\frac{z_1}{q^s z_2}\right) \\ & \quad + \frac{q^{\frac{r+s}{2}}}{1 - q^{r+s}} \delta_{il} \delta_{jk} \left( \delta\left(\frac{z_2}{q^r z_1}\right) - \delta\left(\frac{z_1}{q^s z_2}\right) \right) c. \end{aligned}$$

Comparing the above two identities with the identity (1.19), we derive the following result which was given in [G1].

**Corollary 4.7.** *Let  $M \geq 2$ , and  $G = \langle q \rangle$  be the group generated by  $q \neq 0$  and  $q$  is not a root of unity. Then the Lie algebra  $\mathcal{G}(G, M)$  of operators, acting on  $V_M$ , gives a representation of the Lie algebra  $\widehat{gl}_M(\mathbb{C}_q)$ . The representation is given by the mapping*

$$\begin{aligned} E_{ij} \otimes t_0^m t_1^r &\mapsto \begin{cases} x_{ij}(m, 1, q^r) + \frac{q^{\frac{r}{2}}}{1 - q^r} \delta_{ij} \delta_{m0} c & \text{if } r \neq 0 \\ x_{ij}(m, 1, 1) & \text{if } r = 0 \end{cases} \\ c_0 &\mapsto c, \quad c_1 \mapsto 0 \end{aligned}$$

for  $m, r \in \mathbb{Z}$ .

**Remark 4.8.** The representation of the Lie algebra  $\widehat{gl}_M(\mathbb{C}_Q)$  given in (4.7) is called the homogeneous realization. This is because of the fact that we are using the homogeneous gradation. Moreover the algebra  $\mathcal{G}(G, M)$  contains a subalgebra of  $\mathcal{G}(\langle 1 \rangle, M)$  which is generated by the operators  $x_{ij}(m, 1, 1)$  for  $1 \leq i, j \leq M$  and  $m \in \mathbb{Z}$ , and it is clear that this subalgebra is nothing but the affine algebra  $\widehat{gl}_M(\mathbb{C})$  in the homogeneous picture.

Similarly, we may have the principal realization of the Lie algebra  $\widehat{gl}_N(\mathbb{C}_Q)$ . For this purpose, we choose the group  $G = \langle \xi, q \rangle$  where  $q \neq 0$  is not a root of unity and  $\xi$  is a  $N$ -th primitive root of unity. Let  $M = 1$ . Then the Lie algebra  $\mathcal{G}(G, M)$  is generated by the coefficients of the vertex operators of the form  $X_{11}(\xi^{i-1}, \xi^{-1} q^r, z)$  for all  $r \in \mathbb{Z}$  and  $0 \leq i \leq N - 1$ . From Theorem 3.3 we have

(i) if  $r + s = 0$  and  $\overline{i + j} = 0 \pmod{N}$ , then

$$\begin{aligned} & [X_{11}(\xi^{i-1}, \xi^{-1} q^r, z_1), X_{11}(\xi^{j-1}, \xi^{-1} q^s, z_2)] \\ &= X_{11}(\xi^{i+j-1}, \xi^{-1} q^{r+s}, \xi^{-j} z_1) \delta\left(\frac{\xi^j z_2}{q^r z_1}\right) \\ & \quad - X_{11}(\xi^{i+j-1}, \xi^{-1} q^{r+s}, \xi^{-i} z_2) \delta\left(\frac{\xi^i z_1}{q^s z_2}\right) + (D\delta)\left(\frac{\xi^j z_2}{q^r z_1}\right) c, \end{aligned}$$

(ii) if  $r + s \neq 0$  or  $\overline{i+j} \neq 0(\text{mod } N)$ , then

$$\begin{aligned} & [X_{11}(\xi^{i-1}, \xi^{-1}q^r, z_1), X_{11}(\xi^{j-1}, \xi^{-1}q^s, z_2)] \\ &= X_{11}(\xi^{i+j-1}, \xi^{-1}q^{r+s}, \xi^{-j}z_1) \delta\left(\frac{\xi^j z_2}{q^r z_1}\right) - X_{11}(\xi^{i+j-1}, \xi^{-1}q^{r+s}, \xi^{-i}z_2) \delta\left(\frac{\xi^i z_1}{q^s z_2}\right) \\ &+ \frac{e^{\frac{i+j}{2}Ln\xi} q^{\frac{r+s}{2}}}{\xi^{i+j} - q^{r+s}} \left( \delta\left(\frac{\xi^j z_2}{q^r z_1}\right) - \delta\left(\frac{\xi^i z_1}{q^s z_2}\right) \right) c. \end{aligned}$$

Thus if we write

$$\bar{X}_{11}(\xi^{i-1}, \xi^{-1}q^r, z) = \begin{cases} X_{11}(\xi^{-1}, \xi^{-1}, z) & \text{if } r = 0, i = 0(\text{mod } N) \\ X_{11}(\xi^{i-1}, \xi^{-1}q^r, z) + \frac{e^{\frac{i}{2}Ln\xi} q^{\frac{r}{2}}}{\xi^i - q^r} c & \text{otherwise} \end{cases}$$

for  $r \in \mathbb{Z}$  and  $0 \leq i \leq N-1$ , then the above two identities can be written as one identity

$$\begin{aligned} (4.9) \quad & [\bar{X}_{11}(\xi^{i-1}, \xi^{-1}q^r, z_1), \bar{X}_{11}(\xi^{j-1}, \xi^{-1}q^s, z_2)] \\ &= \bar{X}_{11}(\xi^{i+j-1}, \xi^{-1}q^{r+s}, \xi^{-j}z_1) \delta\left(\frac{\xi^j z_2}{q^r z_1}\right) - \bar{X}_{11}(\xi^{i+j-1}, \xi^{-1}q^{r+s}, \xi^{-i}z_2) \delta\left(\frac{\xi^i z_1}{q^s z_2}\right) \\ &+ \delta_{r+s,0} \delta_{\overline{i+j},0} (D\delta)\left(\frac{\xi^j z_2}{q^r z_1}\right) c. \end{aligned}$$

Comparing this with the identity (1.20), we get the following result which was given in [BS] for the  $N = 2$  case and in [G2] for arbitrary  $N$ .

**Corollary 4.10.** *Let  $M = 1$  and  $G = \langle \xi, q \rangle$  be an admissible subgroup of  $\mathbb{C}^\times$  generated by  $q$  with  $q \neq 0$  not a root of unity, and  $\xi$  is a  $N$ -th primitive root of unity for  $N \geq 2$ . Then the Lie algebra  $\mathcal{G}(G, M)$  of operators, acting on  $V_1$ , gives a representation of the algebra  $\widehat{gl_N}(\mathbb{C}_{q^N})$ . The representation is given by the mapping*

$$\begin{aligned} F^i E^m \otimes t_0^m t_1^r &\mapsto \begin{cases} x_{11}(m, \xi^{i-1}, \xi^{-1}q^r) + \delta_{m,0} \frac{e^{\frac{i}{2}Ln\xi} q^{\frac{r}{2}}}{\xi^i - q^r} c & \text{if } i \neq 0(\text{mod } N) \text{ or } r \neq 0 \\ x_{11}(m, \xi^{-1}, \xi^{-1}) & \text{if } i = 0(\text{mod } N) \text{ and } r = 0 \end{cases} \\ c_0 &\mapsto \frac{c}{N}, \quad c_1 \mapsto 0 \end{aligned}$$

for  $r \in \mathbb{Z}$  and  $0 \leq i \leq N-1$ .

**Remark 4.11.** In general, let  $M, N \geq 2$  be integers,  $G = \langle \xi, q_1, \dots, q_\nu \rangle$  an admissible subgroup of  $\mathbb{C}^\times$  with finitely many generators, where  $q_1, \dots, q_\nu$  are the free generators of  $G$  and  $\xi$  is an  $N$ -th root of unity. Then the Lie algebra  $\mathcal{G}(G, M)$  of operators, acting on the

Fock space  $V_M$ , gives a representation to the Lie algebra  $\widehat{gl_{MN}}(\mathbb{C}_Q)$  where the quantum torus  $\mathbb{C}_Q = \mathbb{C}_Q[t_0^{\pm 1}, t_1^{\pm 1}, \dots, t_\nu^{\pm 1}]$  is determined by the matrix  $Q = (q_{ij})_{(\nu+1) \times (\nu+1)}$  with  $q_{i0} = q_i^N, q_{0i} = q_i^{-N}$  for  $1 \leq i \leq \nu$ , and  $q_{ij} = 1$  for all other values of  $i, j$ .

In particular, if  $\nu = 1$ , that is  $G = \langle \xi, q \rangle$ , then the Lie algebra  $\mathcal{G}(G, M)$  is generated by the coefficient operators of the vertex operators  $X_{ij}(\xi^{k-1}, \xi^{-1}q^r, z)$  for  $1 \leq i, j \leq M$ ,  $1 \leq k \leq N-1$  and  $r \in \mathbb{Z}$ . Moreover, from Theorem 3.3, we have

(i) if  $\overline{k+k'} \neq 0$  or  $r+s \neq 0$ , then

$$\begin{aligned}
& [X_{ij}(\xi^{k-1}, \xi^{-1}q^r, z_1), X_{i'j'}(\xi^{k'-1}, \xi^{-1}q^{r'}, z_2)] \\
&= \delta_{ji'} X_{ij'}(\xi^{k-1}, \xi^{-1-k'}q^{r+r'}, z_1) \delta\left(\frac{\xi^{k'}z_2}{q^r z_1}\right) \\
&\quad - \delta_{j'i} X_{i'j}(\xi^{k'-1}, \xi^{-1-k}q^{r+r'}, z_2) \delta\left(\frac{\xi^k z_1}{q^{r'} z_2}\right) \\
&\quad + \frac{e^{\frac{k+k'}{2}Ln\xi} q^{\frac{r+r'}{2}}}{\xi^{k+k'} - q^{r+r'}} \delta_{ji'} \delta_{j'i} \left(\delta\left(\frac{\xi^{k'}z_2}{q^r z_1}\right) - \delta\left(\frac{\xi^k z_1}{q^{r'} z_2}\right)\right) c \\
&= \delta_{ji'} X_{ij'}(\xi^{k+k'-1}, \xi^{-1}q^{r+r'}, \xi^{-k'}z_1) \delta\left(\frac{\xi^{k'}z_2}{q^r z_1}\right) \\
&\quad - \delta_{j'i} X_{i'j}(\xi^{k+k'-1}, \xi^{-1}q^{r+r'}, \xi^{-k}z_2) \delta\left(\frac{\xi^k z_1}{q^{r'} z_2}\right) \\
&\quad + \frac{e^{\frac{k+k'}{2}Ln\xi} q^{\frac{r+r'}{2}}}{\xi^{k+k'} - q^{r+r'}} \delta_{ji'} \delta_{j'i} \left(\delta\left(\frac{\xi^{k'}z_2}{q^r z_1}\right) - \delta\left(\frac{\xi^k z_1}{q^{r'} z_2}\right)\right) c,
\end{aligned}$$

and

(ii) if  $\overline{k+k'} = 0$  and  $r+s=0$ , then

$$\begin{aligned}
& [X_{ij}(\xi^{k-1}, \xi^{-1}q^r, z_1), X_{i'j'}(\xi^{k'-1}, \xi^{-1}q^{r'}, z_2)] \\
&= \delta_{ji'} X_{ij'}(\xi^{k-1}, \xi^{-1-k'}q^{r+r'}, z_1) \delta\left(\frac{\xi^{k'}z_2}{q^r z_1}\right) \\
&\quad - \delta_{j'i} X_{i'j}(\xi^{k'-1}, \xi^{-1-k}q^{r+r'}, z_2) \delta\left(\frac{\xi^k z_1}{q^{r'} z_2}\right) + \delta_{ji'} \delta_{j'i} (D\delta)\left(\frac{\xi^{k'}z_2}{q^r z_1}\right) c \\
&= \delta_{ji'} X_{ij'}(\xi^{-1}, \xi^{-1}, \xi^{-k'}z_1) \delta\left(\frac{\xi^{k'}z_2}{q^r z_1}\right) \\
&\quad - \delta_{j'i} X_{i'j}(\xi^{-1}, \xi^{-1}, \xi^{-k}z_2) \delta\left(\frac{\xi^k z_1}{q^{r'} z_2}\right) + \delta_{ji'} \delta_{j'i} (D\delta)\left(\frac{\xi^{k'}z_2}{q^r z_1}\right) c.
\end{aligned}$$

Set

$$\overline{X}_{ij}(\xi^{k-1}, \xi^{-1}q^r, z)$$

$$= \begin{cases} X_{ij}(\xi^{k-1}, \xi^{-1}q^r, z) + \delta_{ij} \frac{e^{\frac{k}{2}Ln\xi} q^{\frac{r}{2}}}{\xi^k - q^r} c & \text{if } k \neq 0 \pmod{N} \text{ or } r \neq 0 \\ X_{ij}(\xi^{-1}, \xi^{-1}, z) & \text{if } k = 0 \pmod{N} \text{ and } r = 0 \end{cases}$$

Then we have

$$\begin{aligned} & [\overline{X}_{ij}(\xi^{k-1}, \xi^{-1}q^r, z_1), \overline{X}_{i'j'}(\xi^{k'-1}, \xi^{-1}q^{r'}, z_2)] \\ &= \delta_{ji'} \overline{X}_{ij'}(\xi^{k+k'-1}, \xi^{-1}q^{r+r'}, \xi^{-k'} z_1) \delta\left(\frac{\xi^{k'} z_2}{q^r z_1}\right) \\ & - \delta_{j'i} \overline{X}_{i'j}(\xi^{k+k'-1}, \xi^{-1}q^{r+r'}, \xi^{-k} z_2) \delta\left(\frac{\xi^k z_1}{q^{r'} z_2}\right) + \delta_{ji'} \delta_{j'i} \delta_{k+k',0} \delta_{r+r',0} (D\delta)\left(\frac{\xi^{k'} z_2}{q^r z_1}\right) c. \end{aligned}$$

Comparing this identity with identity (1.21), we get

**Corollary 4.12.** *Let  $M, N \geq 2$ , and let  $G = \langle \xi, q \rangle$  be an admissible subgroup of  $\mathbb{C}^\times$  generated by  $q$  with  $q \neq 0$  not a root of unity, and  $\xi$  an  $N$ -th primitive root of unity. Then the Lie algebra  $\mathcal{G}(G, M)$  of operators, acting on  $V_1$ , gives a representation of the algebra  $\widehat{gl_{MN}}(\mathbb{C}_{q^N})$ , and the representation is given by the mapping*

$$\begin{aligned} & E_{ij} \otimes F^k E^m \otimes t_0^m t_1^r \\ & \mapsto \begin{cases} x_{ij}(m, \xi^{k-1}, \xi^{-1}q^r) + \delta_{m,0} \delta_{ij} \frac{e^{\frac{k}{2}Ln\xi} q^{\frac{r}{2}}}{\xi^k - q^r} c & \text{if } k \neq 0 \pmod{N} \text{ or } r \neq 0 \\ x_{ij}(m, \xi^{-1}, \xi^{-1}) & \text{if } k = 0 \pmod{N} \text{ and } r = 0 \end{cases} \\ & c_0 \mapsto \frac{c}{N}, \quad c_1 \mapsto 0 \end{aligned}$$

for  $r \in \mathbb{Z}$  and  $0 \leq k \leq N-1$ .

The Lie algebra  $\mathcal{G}(G, M)$  given in the previous corollary contains two interesting subalgebras which give representations to the Lie algebras  $\widehat{gl_N}(\mathbb{C}_{q^N})$  and  $\widehat{gl_M}(\mathbb{C}_q)$ . Moreover we will see that these two subalgebras contain subalgebras that give representations to the affine algebras  $\widehat{gl_N}(\mathbb{C})$  of level  $M$  and  $\widehat{gl_M}(\mathbb{C})$  of level  $N$  respectively. Indeed, for  $a, b \in G = \langle \xi, q \rangle$ , let

$$(4.13) \quad Y(a, b, z) = \sum_{k=1}^M X_{kk}(a, b, z)$$

and formally write

$$(4.14) \quad Y(a, b, z) = \sum_{m \in \mathbb{Z}} y(m, a, b) z^{-m}.$$

Let  $\mathcal{L}_1$  be the Lie algebra generated by all of the coefficients of  $Y(a, b, z)$  for  $a, b \in G$ . We note that

$$(4.15) \quad Y(\xi^i q^r, \xi^j q^s, z) = Y(\xi^{i-j-1}, \xi^{-1} q^{s-r}, \xi^{j+1} q^r z),$$

so  $\mathcal{L}_1$  is indeed generated by the coefficients of the vertex operators with the form  $Y(\xi^{i-1}, \xi^{-1} q^r, z)$  for  $r \in \mathbb{Z}$  and  $0 \leq i \leq N-1$ . Moreover, applying Theorem 3.3, we have, if  $r+s \neq 0$  or  $\overline{i+j} \neq 0(\text{mod } N)$

$$(4.16) \quad \begin{aligned} & [Y(\xi^{i-1}, \xi^{-1} q^r, z_1), Y(\xi^{j-1}, \xi^{-1} q^s, z_2)] \\ &= \sum_{k=1}^M [X_{kk}(\xi^{i-1}, \xi^{-1} q^r, z_1), X_{kk}(\xi^{j-1}, \xi^{-1} q^s, z_2)] \\ &= \sum_{k=1}^M \left( X_{kk}(\xi^{i+j-1}, \xi^{-1} q^{r+s}, \xi^{-j} z_1) \delta\left(\frac{\xi^j z_2}{q^r z_1}\right) - X_{kk}(\xi^{i+j-1}, \xi^{-1} q^{r+s}, \xi^{-j} z_2) \delta\left(\frac{\xi^i z_1}{q^s z_2}\right) \right. \\ & \quad \left. + \frac{e^{\frac{i+j}{2} L n \xi q^{\frac{r+s}{2}}}}{\xi^{i+j} - q^{r+s}} \left( \delta\left(\frac{\xi^j z_2}{q^r z_1}\right) - \delta\left(\frac{\xi^i z_1}{q^s z_2}\right) \right) \right) \\ &= Y(\xi^{i+j-1}, \xi^{-1} q^{r+s}, \xi^{-j} z_1) \delta\left(\frac{\xi^j z_2}{q^r z_1}\right) - Y(\xi^{i+j-1}, \xi^{-1} q^{r+s}, \xi^{-j} z_2) \delta\left(\frac{\xi^i z_1}{q^s z_2}\right) \\ & \quad + \frac{e^{\frac{i+j}{2} L n \xi q^{\frac{r+s}{2}}}}{\xi^{i+j} - q^{r+s}} \left( \delta\left(\frac{\xi^j z_2}{q^r z_1}\right) - \delta\left(\frac{\xi^i z_1}{q^s z_2}\right) \right), \end{aligned}$$

while if  $r+s=0$  and  $\overline{i+j}=0(\text{mod } N)$ , then

$$(4.17) \quad \begin{aligned} & [Y(\xi^{i-1}, \xi^{-1} q^r, z_1), Y(\xi^{j-1}, \xi^{-1} q^s, z_2)] \\ &= \sum_{k=1}^M [X_{kk}(\xi^{i-1}, \xi^{-1} q^r, z_1), X_{kk}(\xi^{j-1}, \xi^{-1} q^s, z_2)] \\ &= \sum_{k=1}^M \left( X_{kk}(\xi^{-1}, \xi^{-1}, \xi^{-j} z_1) \delta\left(\frac{\xi^j z_2}{q^r z_1}\right) - X_{kk}(\xi^{-1}, \xi^{-1}, \xi^{-j} z_2) \delta\left(\frac{\xi^i z_1}{q^s z_2}\right) + (D\delta)\left(\frac{\xi^j z_2}{q^r z_1}\right) \right) \\ &= Y(\xi^{-1}, \xi^{-1}, \xi^{-j} z_1) \delta\left(\frac{\xi^j z_2}{q^r z_1}\right) - Y(\xi^{-1}, \xi^{-1}, \xi^{-j} z_2) \delta\left(\frac{\xi^i z_1}{q^s z_2}\right) + M(D\delta)\left(\frac{\xi^j z_2}{q^r z_1}\right). \end{aligned}$$

Therefore, if we define

$$\bar{Y}(\xi^{i-1}, \xi^{-1}q^r, z) = \begin{cases} Y(\xi^{i-1}, \xi^{-1}q^r, z) + M \frac{e^{\frac{i}{2}Ln\xi} q^{\frac{r}{2}}}{\xi^i - q^r} c & \text{if } r \neq 0 \text{ or } \bar{i} \neq 0 \\ Y(\xi^{-1}, \xi^{-1}, z) & \text{if } r = 0 \text{ and } \bar{i} = 0, \end{cases}$$

then we can rewrite the two identities (4.16) and (4.17) into just one identity

$$(4.18) \quad [\bar{Y}(\xi^{i-1}, \xi^{-1}q^r, z_1), \bar{Y}(\xi^{j-1}, \xi^{-1}q^s, z_2)] \\ = \bar{Y}(\xi^{-1}, \xi^{-1}, \xi^{-j}z_1) \delta\left(\frac{\xi^j z_2}{q^r z_1}\right) \bar{Y}(\xi^{-1}, \xi^{-1}, \xi^{-j}z_2) \delta\left(\frac{\xi^i z_1}{q^s z_2}\right) + M(D\delta)\left(\frac{\xi^j z_2}{q^r z_1}\right) c.$$

Therefore we have the following result

**Proposition 4.19.** *The Lie algebra  $\mathcal{L}_1$  of operators acting on  $V_M$  gives a representation of the Lie algebra  $\widehat{gl}_N(\mathbb{C}_{q^N})$ , and the representation is given by the mapping*

$$F^i E^m \otimes t_0^m t_1^r \mapsto \begin{cases} y(m, \xi^{i-1}, \xi^{-1}q^r) + M \frac{e^{\frac{i}{2}Ln\xi} q^{\frac{r}{2}}}{\xi^i - q^r} \delta_{m,0} c & \text{if } i \neq 0 \pmod{N} \text{ or } r \neq 0 \\ y(m, 1, 1) & \text{if } i = 0 \pmod{N} \text{ and } r = 0 \end{cases} \\ c_0 \mapsto Mc, \quad c_1 \mapsto 0$$

for  $r \in \mathbb{Z}$  and  $0 \leq i \leq N-1$ .

Recall from (4.7), the Fock space  $V_M$  affords a representation of the Lie algebra  $\mathcal{G}(< q >, M) \subset \mathcal{G}(< \xi, q >, M)$ , where  $\xi, q$  are given in (4.10), and

$$\mathcal{G}(< q >, M) = \text{span} \{c \text{ and } x_{ij}(m, 1, q^r) \mid \text{for } m, r \in \mathbb{Z}, 1 \leq i, j \leq M\}.$$

Now we define a subalgebra of  $\mathcal{G}(< q >, M) \subset \mathcal{G}(< \xi, q >, M)$

$$\mathcal{L}_2 = \text{span}\{c \text{ and } x_{ij}(Nm, 1, q^r) \mid \text{for } m, r \in \mathbb{Z}, 1 \leq i, j \leq M\}.$$

Then we have

**Proposition 4.20.**  *$\mathcal{L}_2$  forms a Lie subalgebra of  $\mathcal{G}(< q >, M)$ , and  $\mathcal{L}_2$  is also isomorphic to  $\mathcal{G}(< q >, M)$  via the isomorphism given by*

$$x_{ij}(m, 1, q^r) \mapsto x_{ij}(Nm, 1, q^r), \quad c \mapsto Nc.$$

Therefore  $\mathcal{L}_2$  gives a representation of  $\widehat{gl}_N(\mathbb{C}_{q^N})$ .

**Proposition 4.21.** For  $m, n, r, s \in \mathbb{Z}$  and  $i \not\equiv 0 \pmod{N}$ ,  $1 \leq k \neq l \leq M$ , we have

$$(4.22) \quad \begin{aligned} & [y(m, \xi^{i-1}, \xi^{-1}q^r), x_{kl}(Nn, 1, q^s)] \\ &= (q^{rNn} - q^{sm})x_{kl}(m + Nn, \xi^{i-1}, \xi^{-1}q^{r+s}). \end{aligned}$$

*Proof.* We apply Theorem 3.3 to obtain

$$\begin{aligned} & [Y(\xi^{i-1}, \xi^{-1}q^r, z_1), X_{kl}(1, q^s, z_2)] = [\sum_{j=1}^M X_{jj}(\xi^{i-1}, \xi^{-1}q^r, z_1), X_{kl}(1, q^s, z_2)] \\ &= \sum_{j=1}^M \{X_{jl}(\xi^{i-1}, \xi^{-1}q^{r+s}, z_1)\delta_{jk}\delta(\frac{z_2}{\xi^{-1}q^r z_1}) - X_{kj}(1, \xi^{-i}q^{r+s}, z_2)\delta_{jl}\delta(\frac{\xi^{i-1}z_1}{q^s z_2})\} \\ &= X_{kl}(\xi^{i-1}, \xi^{-1}q^{r+s}, z_1)\delta(\frac{z_2}{\xi^{-1}q^r z_1}) - X_{kl}(1, \xi^{-i}q^{r+s}, z_2)\delta(\frac{\xi^{i-1}z_1}{q^s z_2}) \\ &= X_{kl}(\xi^{i-1}, \xi^{-1}q^{r+s}, z_1)\delta(\frac{z_2}{\xi^{-1}q^r z_1}) - X_{kl}(\xi^{i-1}, \xi^{-1}q^{r+s}, q^{-s}z_1)\delta(\frac{\xi^{i-1}z_1}{q^s z_2}). \end{aligned}$$

This then gives

$$[y(m, \xi^{i-1}, \xi^{-1}q^r), x_{kl}(n, 1, q^s)] = \xi^{-n}(q^{rn} - q^{sm}\xi^{in})x_{kl}(m + n, \xi^{i-1}, \xi^{-1}q^{r+s}).$$

which immediately implies (4.22).  $\square$

**Remark 4.23** Let  $\mathcal{G}_i \subset \mathcal{L}_i \subset \mathcal{G}(< \xi, q >, M)$ ,  $i = 1, 2$ , be such that

$$\begin{aligned} \mathcal{G}_1 &= \text{span}\{c \text{ and } y(m, \xi^{i-1}, \xi^{-1}) \mid \text{ for } m \in \mathbb{Z}, 0 \leq i \leq N-1\}, \\ \mathcal{G}_2 &= \text{span}\{c \text{ and } x_{ij}(Nm, 1, 1) \mid \text{ for } m \in \mathbb{Z}, 1 \leq i, j \leq M\}. \end{aligned}$$

Then the two subalgebras  $\mathcal{G}_1, \mathcal{G}_2$  of  $\mathcal{G}(< \xi, q >, M)$  respectively give representations of the affine algebra  $\widehat{gl_N}(\mathbb{C})$  of level  $M$  and  $\widehat{gl_M}(\mathbb{C})$  of level  $N$ . Let  $\mathcal{G}'_i$  be the derived algebras of  $\mathcal{G}_i$ . Then we have the so called dual pair property given in [F1]:  $[\mathcal{G}'_1, \mathcal{G}'_2] = (0)$ . However, clearly, we have  $[\mathcal{L}'_1, \mathcal{L}'_2] \neq (0)$ .

## Acknowledgements

SB and YG are supported by grants from the Natural Sciences and Engineering Research Council of Canada. ST is supported by a grant from the National Natural Science Foundation of China. The authors would like to thank the Fields Institute for its hospitality during the preparation of this work.

## REFERENCES

- [AABGP] B. N. Allison, S. Azam, S. Berman, Y. Gao, A. Pianzola, *Extended affine Lie algebras and their root systems*, Memoir. Amer. Math. Soc. **126** (1997), Number 605.
- [ABGP] B. N. Allison, S. Berman, Y. Gao, A. Pianzola, *A characterization of affine Kac-Moody Lie algebras*, Comm. Math. Phys. **185** (1997), 671–688.
- [BC] S. Berman and B. Cox, *Enveloping algebras and representations of toroidal Lie algebras*, Pacific J. Math **165** (1994), 239–267.
- [BGK] S. Berman, Y. Gao, Y. Krylyuk, *Quantum tori and the structure of elliptic quasi-simple Lie algebras*, J. Funct. Anal **135** (1996), 339–389.
- [BS] S. Berman and J. Szmigielski, *Principal realization for extended affine Lie algebra of type  $sl_2$  with coordinates in a simple quantum torus with two variables*, Cont.Math. **248** (1999), 39–67.
- [F1] I. B. Frenkel, *Representations of affine Lie algebras, Hecke modular forms and Korteweg-De Vries type equations*, Lecture notes in Math. **933** (1982), 71–110.
- [F2] I. B. Frenkel, *Representations of Kac-Moody algebras and dual resonance models*, Lectures in Appl. Math. **21** (1985), 325–353.
- [F3] I.B. Frenkel, *Spinor representations of affine Lie algebras*, Proc. Nat. Acad. Sci. USA **77** (1980), 6303–6306.
- [F4] I.B. Frenkel, *Two constructions of affine Lie algebra representations and Boson–Fermion correspondence in quantum field theory*, J. Funct. Anal. **44** (1981), 259–327.
- [FJW] I. B. Frenkel, N. Jing and W. Wang, *Vertex representations via finite groups and the McKay correspondence*, IMRN **4** (2000), 195–222.
- [FK] I. B. Frenkel and V. G. Kac, *Representations of affine Lie algebras and dual resonance models*, Invent. Math. **62** (1980), 23–66.
- [FLM] I. B. Frenkel, J. Lepowsky and A. Meurman, *Vertex Operator Algebras and the Monster*, Academic Press, 1989.
- [G1] Y. Gao, *Vertex operators arising from the homogeneous realization for  $\widehat{gl}_N$* , Comm. Math. Phys. **211** (2000), 745–777.
- [G2] Y. Gao, *Representations of extended affine Lie algebras coordinatized by certain quantum tori*, Compositio Mathematica **123** (2000), 1–25.
- [G3] Y. Gao, *Fermionic and bosonic representations of the extended affine Lie algebra  $\widetilde{\mathfrak{gl}_N(\mathbb{C}_q)}$* , Submitted.
- [G-KK] M. Golenishcheva-Kutuzova and V. G. Kac,  *$\Gamma$ -conformal algebras*, J. Math. Phys. **39** (1998), 2290–2305.
- [G-KL] M. Golenishcheva-Kutuzova and D. Lebedev, *Vertex operator representation of some quantum tori Lie algebras*, Comm. Math. Phys. **148** (1992), 403–416.
- [J] N. H. Jing, *Twisted vertex representations of quantum affine algebras*, Invent. Math. **102** (1990), 663–690.
- [H-KT] R. Høegh-Krohn and B. Torresani, *Classification and construction of quasi-simple Lie algebras*, J. Funct. Anal. **89** (1990), 106–136.
- [JK] H. P. Jakobsen and V. G. Kac, *A new class of unitarizable highest weight representations of infinite-dimensional Lie algebras. II*, J. Funct. Anal. **82** (1989), 69–90.
- [K] V. G. Kac, *Infinite dimensional Lie algebras*, third edition, Cambridge Univ. Press, 1990.
- [KP] V. Kac and D. Peterson, *Spin and wedge representations of infinite-dimensional Lie algebras and groups*, Proc. Nat. Acad. Sci. USA **78** (1981), 3308–3312.
- [KKLW] V. G. Kac, D. A. Kazhdan, J. Lepowsky and R. L. Wilson, *Realization of the basic representations of the Euclidean Lie algebras*, Advances in Math. **42** (1981), 83–112.



- [LW] J. Lepowsky and R. L. Wilson, *Construction of the affine Lie algebra  $A_1^{(1)}$* , Comm. Math. Phys. **62** (1978), 43–53.
- [M] Y. I. Manin, *Topics in noncommutative geometry*, Princeton University Press, 1991.
- [MRY] R. V. Moody, S. E. Rao and T. Yokonuma, *Toroidal Lie algebras and vertex representations*, Geom. Ded. **35** (1990), 283–307.
- [S] G. Segal, *Unitary representations of some infinite-dimensional groups*, Comm. Math. Phys. **80** (1981), 301–342.
- [T1] S. Tan, *TKK algebras and vertex operator representations*, J. Algebra **211** (1999), 298–342.
- [T2] S. Tan, *Vertex operator representations for toroidal Lie algebras of type  $B_l$* , Comm. Algebra **27** (1999), 3593–3618.
- [VV] M. Varagnolo and E. Vasserot, *Double-loop algebras and the Fock space*, Invent. Math. **133** (1998), 133–159.
- [W] M. Wakimoto, *Extended affine Lie algebras and a certain series of Hermitian representations*, Preprint (1985).
- [Y] H. Yamada, *Extended affine Lie algebras and their vertex representations*, Publ. RIMS, Kyoto U. **25** (1989), 587–603.